


(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 072 592 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication
31.01.2001 Bulletin 2001/05

(51) Int Cl⁷ C07D 211/14, C07D 211/22,
C07D 211/32, C07D 211/34,
A61K 31/445, A61P 17/04

(21) Application number. 00304219.9

(22) Date of filing 18.05.2000

BEST AVAILABLE COPY

(84) Designated Contracting States
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States
AL LT LV MK RO SI

(30) Priority 28.05.1999 GB 9912411

(71) Applicants
• PFIZER INC.
New York, N.Y. 10017-5755 (US)
Designated Contracting States
BE CH DE DK ES FI FR GR IE IT LI LU NL PT AT CY
• Pfizer Limited
Sandwich Kent CT13 9NJ (GB)
Designated Contracting States
GB


(72) Inventors
• Armer, Richard Edward
Newhouse, Scotland ML1 5SH (GB)
• Gethin, David Morris
Sandwich, Kent CT13 9NJ (GB)
• Gibson, Stephen Paul
Sandwich, Kent CT13 9NJ (GB)
• Tommasini, Ivan
Sandwich, Kent CT13 9NJ (GB)

(74) Representative Eddowes, Simon et al
Urquhart-Dykes & Lord,
30 Welbeck Street
London W1G 8ER (GB)

(54) 4-arylpiperidine derivatives for the treatment of pruritus

(57) There is provided a compound of formula I,

wherein A, D, R¹, R², R³, X and n have meanings given in the description, which are useful in the prophylaxis and in the treatment of diseases mediated by opiate receptors such as pruritus



EP 1 072 592 A2

Description

[0001] This invention relates to pharmaceutically useful compounds, in particular compounds that bind to opiate receptors (e.g. μ , κ and δ opiate receptors)

5 [0002] Compounds that bind to such receptors are likely to be useful in the treatment of diseases mediated by opiate receptors, for example irritable bowel syndrome, constipation, nausea, vomiting, and pruritic dermatoses, such as allergic dermatitis and atopy in animals and humans. Compounds that bind to opiate receptors have also been indicated in the treatment of eating disorders, opiate overdoses, depression, smoking and alcohol addiction, sexual dysfunction, shock, stroke, spinal damage and head trauma.

10 [0003] There is a particular need for an improved treatment of itching. Itching, or pruritus, is a common dermatological symptom that can give rise to considerable distress in both humans and animals. Pruritus is often associated with inflammatory skin diseases which may be caused by hypersensitivity reactions, including reactions to insect bites, such as flea bites, and to environmental allergens, such as house dust mite or pollen, by bacterial and fungal infections of the skin or by ectoparasite infections.

15 [0004] Existing treatments that have been employed in the treatment of pruritus include the use of corticosteroids and antihistamines. However, both of these treatments are known to have undesirable side effects. Other therapies that have been employed include the use of essential fatty acid dietary supplements, though these have the disadvantages of being slow to act, and of offering only limited efficacy against allergic dermatitis. A variety of emollients such as soft paraffin, glycerine and lanolin are also employed, but with limited success.

20 [0005] Thus, there is a continuing need for alternative and/or improved treatments of pruritus.

[0006] Certain 4-arylpiperidine-based compounds are disclosed in *inter alia* European patent applications EP 267339, EP 506468, EP 506478 and *J Med Chem* 1993, 36, 2833-2850 as opiate antagonists. In addition, International Patent Application WO 95/15327 discloses azabicycloalkane derivatives useful as neuroleptic agents.

[0007] According to the invention there is provided compounds of formula I

25

30

35

40

wherein

45

50

55

A represents a single bond, C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH.

D represents H, OH, CN, $N(R^4)(R^5)$, $N(H)R^6$, $C(O)N(R^4)(R^5)$, $C(O)OR^7$, $C(O)R^8$, $C(=NR^{9a})R^8$, or $C(=NOR^{9b})R^8$, provided that when A represents C_{2-4} alkenylene or C_{2-4} alkynylene, and D represents OH, $N(R^4)(R^5)$ or $N(H)R^6$, then D is not directly attached to an unsaturated carbon atom;

and provided that when A represents a single bond, then D does not represent H, OH, $N(R^4)(R^5)$ or $N(H)R^6$.

R^4 and R^5 independently represent H, C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), or R^4 and R^5 together with the N-atom to which they are attached, form a 4- to 7-membered heterocyclic ring, which ring optionally contains one or more additional heteroatoms selected from oxygen, nitrogen and sulfur and which ring is optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, OH, =O, nitro, amino or halo.

R^6 represents $C(O)R^{10a}$, $C(O)OR^{10b}$ or $S(O)_2R^{10c}$.

R^{10a} to R^{10c} independently represent C_{1-4} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl (which four groups are all optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which

latter two groups are optionally substituted by one or more halo atoms)), or R^{10a} represents H;
 R^7 and R^8 independently represent H, C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl or C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms);

R^{9a} and R^{9b} independently represent C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), or R^{9b} represents H;

R^1 and R^2 are each independently H or C_{1-4} alkyl.

R^3 represents aryl (optionally substituted by one or more substituents selected from OH, nitro, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms) and $-N(R^{11a})(R^{11b})$), C_{1-10} alkyl, C_{3-10} alkenyl or C_{3-10} alkynyl wherein said alkyl, alkenyl or alkynyl groups are optionally substituted and/or terminated by one or more substituents selected from OR^{11c} , $S(O)_pR^{11d}$, CN, halo, C_{1-6} alkoxy carbonyl, C_{2-6} alkanoyl, C_{2-6} alkanoyloxy, C_{3-8} cycloalkyl, C_{4-9} cycloalkanoyl, $N(R^{12a})S(O)_2R^{13}$, Het^1 , aryl, adamantyl (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)) or $-W-A^1-N(R^{12b})(R^{12c})$.

p is 0, 1 or 2,

W represents a single bond, $C(O)$ or $S(O)_q$,

A^1 represents a single bond or C_{1-10} alkylene;

provided that when both W and A^1 represent single bonds, then the group $-N(R^{12b})(R^{12c})$ is not directly attached to an unsaturated carbon atom;

q is 0, 1 or 2;

R^{11a} to R^{11d} each independently represent H, C_{1-10} alkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{3-8} cycloalkyl, C_{1-4} alkylphenyl, aryl (which latter six groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)) or Het^2 ;

provided that R^{11d} does not represent H when p represents 1 or 2,

R^{12a} to R^{12c} each independently represent H, C_{1-10} alkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{3-8} cycloalkyl, C_{1-4} alkylphenyl, aryl (which latter six groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), Het^3 , or R^{12b} and R^{12c} together represent unbranched C_{2-4} alkylene which alkylene group is optionally interrupted by O, S and/or an $N(R^{14})$ group and is optionally substituted by one or more C_{1-4} alkyl groups;

R^{13} represents C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{1-4} alkylphenyl or aryl, which four groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, OH, nitro, amino or halo,

R^{14} represents H, C_{1-6} alkyl, C_{3-8} cycloalkyl, $A^2-(C_{3-8}$ cycloalkyl) or A^2 -aryl,

A^2 represents C_{1-6} alkylene,

Het^1 , Het^2 and Het^3 independently represent 3- to 8-membered heterocyclic groups, which groups contain at least one heteroatom selected from oxygen, sulfur and/or nitrogen, which groups are optionally fused to a benzene ring, and which groups are optionally substituted in the heterocyclic and/or fused benzene ring part by one or more substituents selected from OH, =O, nitro, amino, halo, CN, aryl, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms);

X is H, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms),

n is 0, 1 or 2,

or pharmaceutically, or veterinarily, acceptable derivatives thereof,

which compounds are referred to together hereinafter as "the compounds of the invention"

[0008] In the definitions used herein, alkyl, alkylene, alkoxy, alkoxy carbonyl, alkanoyl, alkanoyloxy, alkenyl, alkynyl and the alkyl parts of alkylphenyl and aryl alkoxy groups may, when there is a sufficient number of carbon atoms, be straight or branched-chain and/or optionally interrupted by one or more oxygen and/or sulfur atom(s). The term halo includes fluoro, chloro, bromo or iodo. The term "aryl" includes optionally substituted phenyl, naphthyl and the like, and "aryloxy" includes optionally substituted phenoxy and naphthyloxy and the like. Unless otherwise specified, aryl and aryloxy groups are optionally substituted by one or more (e.g. one to three) substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy carbonyl and C_{1-5} alkanoyl (which latter four groups are optionally substituted by one or more halo atoms).

[0009] The heterocyclic rings that Het^1 , Het^2 and Het^3 represent and that $N(R^4)(R^5)$ may represent, may be fully saturated, partially unsaturated and/or wholly or partially aromatic in character.

[0010] For the avoidance of doubt when heterocyclic groups (i.e. Het¹, Het², Het³ and some definitions of N(R⁴)(R⁵)) are at least part-saturated, possible points of substitution include the atom (e.g. the carbon atom) at the point of attachment of the heterocyclic group to the rest of the molecule. Het (Het¹, Het² and Het³) groups may also be attached to the rest of the molecule via a heteroatom

5 [0011] The piperidine moiety in compounds of formula I may be in N-oxidised form. Sulfur atoms that may interrupt (e.g. alkyl) substituents in compounds of formula I may be present in oxidised form (e.g. as sulfoxides or sulfones). All heterocyclic groups (i.e. Het¹, Het², Het³ and some definitions of N(R⁴)(R⁵)) may also be in N- or S-oxidized forms

[0012] The term "pharmaceutically, or veterinarily, acceptable derivatives" includes non-toxic salts. Salts which may be mentioned include acid addition salts, for example, salts formed with sulfuric, hydrochloric, hydrobromic, phosphoric, hydriodic, sulfamic, organo-sulfonic, citric, carboxylic (e.g. acetic, benzoic, etc.), maleic, malic, succinic, tartaric, cinnamic, ascorbic and related acids, base addition salts, salts formed with bases, for example, the sodium, potassium and C₁₋₄ alkyl ammonium salts

[0013] The compounds of the invention may also be in the form of quaternary ammonium salts e.g. at the piperidine moiety, which salts may be formed by reaction with a variety of alkylating agents, such as an alkyl halide or an ester of sulfuric or an aromatic sulfonic acid

15 [0014] The compounds of the invention may exhibit tautomerism. All tautomeric forms of the compounds of formula I are included within the scope of the invention

[0015] The compounds of the invention contain one or more asymmetric centres and thus they can exist as enantiomers and diastereomers. Diastereoisomers may be separated using conventional techniques e.g. by fractional crystallisation or chromatography. The various stereoisomers may be isolated by separation of a racemic or other mixture of the compounds using conventional techniques e.g. fractional crystallisation or HPLC. The desired optical isomers may be prepared by reaction of the appropriate optically active starting materials under conditions which will not cause racemisation or epimerisation. Alternatively, the desired optical isomers may be prepared by resolution, either by HPLC of the racemate using a suitable chiral support or, where appropriate, by fractional crystallisation of the diastereoisomeric salts formed by reaction of the racemate with a suitable optically active acid or base. The invention includes the use of both the separated individual isomers as well as mixtures of isomers

[0016] Also included within the scope of the invention are radiolabelled derivatives of compounds of formula I which are suitable for biological studies

[0017] Preferred compounds of the invention include those wherein

30 The group A-D is attached in the *meta*- position relative to the piperidine ring.
 R¹ represents C₁₋₂ alkyl.
 R² represents H or C₁₋₂ alkyl.
 R³ represents saturated C₁₋₁₀ (e.g. C₁₋₈) alkyl optionally substituted by one or more substituents selected from OR^{11c}, CN, halo, C₂₋₄ alkanoyl, C₁₋₄ alkoxy carbonyl, N(R^{12a})SO₂R¹³, Het¹, aryl (which latter group is optionally substituted by one or more substituents selected from OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₅ alkanoyl, halo, nitro, amino, CN and CONH₂), or -W-A¹-N(R^{12b})(R^{12c}).
 R^{11c} represents H, C₁₋₆ alkyl or aryl (which latter groups is optionally substituted by one or more substituents selected from OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₅ alkanoyl, halo, nitro, amino, CN and CONH₂);
 40 R^{12a} to R^{12c} independently represent H, C₁₋₄ alkyl, C₁₋₂ alkylphenyl or aryl (which latter three groups are optionally substituted by one or more substituents selected from halo, C₁₋₄ alkyl or C₁₋₄ alkoxy)
 R¹³ represents C₁₋₄ alkyl, C₁₋₂ alkylphenyl or aryl (which three groups are all optionally substituted by one or more substituents selected from halo, C₁₋₄ alkyl or C₁₋₄ alkoxy),
 W represents C(O)
 45 A¹ represents a single bond

[0018] More preferred compounds of the invention include those wherein

50 A represents a single bond, C₁₋₄ alkylene, C₂₋₄ alkenylene or C₂₋₄ alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more OH and/or methyl groups,
 D represents H, OH, CN, N(H)R⁴, N(H)C(O)R^{10a}, N(H)C(O)OR^{10b}, N(H)S(O)₂R^{10c}, C(O)N(R⁴)(R⁵), C(O)OR⁷, C(O)R⁸ or C(=NOH)R⁸,
 R⁴ and R⁵ independently represent H, C₁₋₄ alkyl or C₁₋₃ alkylphenyl (which latter two groups are both optionally substituted by C₁₋₄ alkoxy)
 55 R⁷ and R⁸ independently represent H or C₁₋₄ alkyl
 R^{10a} to R^{10c} independently represent C₁₋₄ alkyl (optionally substituted by one or more halo atoms)
 R¹ represents methyl,
 R² represents H or methyl

R³ represents saturated C₁₋₇ alkyl, optionally substituted by one or more substituents selected from CN, OR^{11c} or phenyl.

R^{11c} represents C₁₋₆ alkyl or phenyl;

X represents halo, particularly fluoro;

n represents 1 or, preferably, 0

[0019] Particularly preferred compounds of the invention include those wherein

A represents a single bond, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -CH(OH)-, -(CH₂)₂-, -CH=CH-, or -C≡C-;

D represents H, OH, CN, NH₂, N(H)CH₃, CHO, CH(=NOH), C(O)CH₃, CO₂CH₃, CO₂H, C(O)NH₂, C(O)N(H)CH₃, C(O)N(H)Et, C(O)N(H)(2-MeOEt), C(O)N(H)*n*-Pr, C(O)N(H)*i*-Pr, C(O)N(H)*n*-Bu, C(O)N(H)*i*-Bu, C(O)N(H)*t*-Bu, C(O)N(H)CH₂Ph, C(O)N(CH₃)₂, C(O)N(Et)₂, N(H)C(O)CH₃, N(H)C(O)OCH₃, N(H)S(O)₂CH₃ or N(H)S(O)₂CF₃,

R¹ and R² represent methyl groups in the mutually *trans* configuration,


R³ represents benzyl, 5-cyanopentyl, *n*-hexyl, 5-methylhexyl, 2-phenoxyethyl or 3-phenylpropyl

[0020] Preferred compounds of the invention include the compounds of the Examples described hereinafter


[0021] According to a further aspect of the invention there is provided processes for the preparation of compounds of the invention, as illustrated below

[0022] The following processes are illustrative of the general synthetic procedures which may be adopted in order to obtain the compounds of the invention

1. Compounds of formula I in which A represents C₂₋₄ alkynylene (in which group the carbon-carbon triple bond is α, β to the benzene ring) which alkynylene group is optionally substituted at the 3- and/or the 4-C (relative to the benzene ring) by one or more substituents defined hereinbefore in respect of A, and/or one of the groups defined hereinbefore in respect of D, or (when D is not attached at the 3- or 4-C) which alkynylene group is substituted at the 2-C (relative to the benzene ring) by CN, C(O)N(R⁴)(R⁵), C(O)OR⁷, C(O)R⁶, C(=NR^{9a})R⁸, or C(=NOR^{9b})R⁸ may be prepared by reaction of a corresponding compound of formula II,




wherein L¹ is a suitable leaving group such as halogen, preferably bromine or iodine, or a sulfonate such as trifluoromethanesulfonate, and R¹, R², R³, X and n are as hereinbefore defined, with a compound of formula III,



wherein M represents (as appropriate) H, a tin-containing moiety (e.g. tributylstannyl), a boron derivative (e.g. a boronic acid), a zinc halide, a magnesium halide or an alkali metal (which latter three groups may be formed *in situ* from the corresponding halide), A³ represents a single bond or C₁₋₂ alkylene (optionally substituted by one or more substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, OH or halo), and D is as hereinbefore defined, provided that when A³ represents a single bond, then D does not represent H, OH, N(R⁴)(R⁵) or N(H)R⁶, wherein R⁴, R⁵ and R⁶ are as hereinbefore defined, for example at between room and reflux temperature in the presence of a suitable catalyst system (e.g. bis(triphenylphosphine)palladium(II) chloride combined with copper(I) iodide) and an appropriate organic base (e.g. triethylamine).

2. Compounds of formula I in which A represents C₂₋₄ alkenylene (in which group the carbon-carbon double bond


is α, β to the benzene ring), which alkenylene group is optionally substituted at the 2-C (relative to the benzene ring) by C_{1-4} alkyl, and also optionally substituted at the 3- and/or 4-C (relative to the benzene ring) by one or more of the substituents defined hereinbefore in respect of A and/or one of the groups defined hereinbefore in respect of D, or which alkenylene group is substituted at the 2-C (relative to the benzene ring) by CN, $C(O)N(R^4)(R^5)$, $C(O)OR^7$, $C(O)R^8$, $C(=NR^{9a})R^8$, or $C(=NOR^{9b})R^8$, may be prepared by reaction of a corresponding compound of formula II, as hereinbefore defined, with a compound of formula IV,



IV

wherein the dashed bond represent optional *cis*- or *trans*- geometry, R^{15} represents H or C_{1-4} alkyl, and A^3 , D and M are as hereinbefore defined. for example at between room temperature and reflux temperature in the presence of a reaction-inert solvent (e.g. 1,4-dioxan or THF), an appropriate catalyst (e.g. tetrakis(triphenylphosphine)palladium(0) or bis(triphenylphosphine)palladium(II) acetate) and either (as appropriate) a suitable source of halide ion (e.g. lithium chloride) or a suitable base (e.g. triethylamine)


3 Compounds of formula I in which A represents a single bond and D represents CN may be prepared by reaction of a compound of formula V,



V

wherein R^1 , R^2 , R^3 , X and n are as hereinbefore defined with an alkali metal cyanide (e.g. potassium cyanide), for example at raised temperature in the presence of a reaction-inert solvent (e.g. N-methylpyrrolidine) and a suitable catalyst (e.g. palladium(II) acetate combined with 1,1'-bis(diphenylphosphino)ferrocene)


Compounds of formula V may be prepared by reaction of a corresponding compound of formula VI,



VI

wherein R^1 , R^2 , R^3 , X and n are as hereinbefore defined, with an appropriate triflating agent (e.g. N-phenyltrifluoromethanesulfonimide), for example at between 0°C and room temperature in the presence of a reaction-inert organic solvent (e.g. dichloromethane) and a suitable base (e.g. triethylamine)

Compounds of formula VI may be prepared by reaction of a corresponding compound of formula VII



VII

in which R^1 , R^2 , X and n are as hereinbefore defined, with a compound of formula VIII,



VIII

wherein R^3 and L^1 are as hereinbefore defined, under conditions that are known to those skilled in the art, which include, for example, alkylation at between room temperature and reflux temperature in the presence of a reaction-inert organic solvent (e.g. *N,N*-dimethylformamide) and a suitable base (e.g. NaHCO_3), and arylation at between room temperature and reflux temperature in the presence of a suitable catalyst system (e.g. tris(dibenzylideneacetone)palladium(0) combined with tri-*o*-tolylphosphine), an appropriate strong base (e.g. sodium *tert*-butoxide) and a reaction-inert solvent (e.g. toluene).

4 Compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents NH_2 (which is attached to a CH_2 group) may be prepared by reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents CN, for example at between room and reflux temperature in the presence of a suitable reducing agent (e.g. lithium aluminium hydride) and an appropriate solvent (e.g. THF).


5 Compounds of formula I in which D represents C(O)NH_2 may be prepared by controlled hydrolysis of a corresponding compound of formula I in which D represents CN, for example by reaction with polyphosphoric acid at between 50 and 150°C.

6. Compounds of formula I in which A represents a single bond and D represents $\text{C(O)-(C}_{1-6}\text{ alkyl)}$ or $\text{C(O)-(C}_{1-4}\text{ alkylphenyl)}$, which alkyl and alkylphenyl groups are both optionally substituted by one or more of the substituents defined hereinbefore in respect of R^8 , may be prepared by hydrolysis of a corresponding compound of formula IX,

5

10

15



20

wherein R^{15} represents C_{1-6} alkyl, R^{16} represents H, C_{1-5} alkyl, phenyl or C_{1-3} alkylphenyl which latter three groups are all optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), the dashed bond indicates optional *cis*- or *trans*- geometry, and R^1 , R^2 , R^3 , X and n are as hereinbefore defined, for example under conditions known to those skilled in the art (e.g. by reaction at between room and reflux temperature with an aqueous solution of a mineral acid)

25

Compounds of formula IX may be prepared by reaction of a compound of formula II, as hereinbefore defined, with a compound of formula X

30



35

wherein the dashed bond indicates optional *cis*- or *trans*- geometry, and R^{15} and R^{16} are as hereinbefore defined, for example at between room temperature and reflux temperature in the presence of an appropriate catalyst (e.g. palladium(II) acetate combined with 1,1'-bis(diphenylphosphino)ferrocene), an organic base (e.g. triethylamine) and an appropriate solvent (e.g. *N,N*-dimethylformamide)

40

7 Compounds of formula I in which D represents $C(O)R^8$ wherein R^8 is as hereinbefore defined provided that it does not represent H, may be prepared by reaction of a corresponding compound of formula I in which D represents CN with an organometallic compound capable of delivering an R^{8a} -containing anion (e.g. an appropriate organolithium or Grignard reagent) wherein R^{8a} is defined as for R^8 above provided that it does not represent H, for example at between -80 and 10°C in the presence of a reaction-inert organic solvent (e.g. tetrahydrofuran)

45

8 Compounds of formula I in which A represents a single bond and D represents $C(O)OR^7$, wherein R^7 is as hereinbefore defined provided that it does not represent H, may be prepared by reaction of a corresponding compound of formula V as hereinbefore defined with carbon monoxide and an alcohol of formula $R^{7a}OH$, wherein R^{7a} is defined as for R^7 above provided that it does not represent H, for example in the presence of a suitable transition-metal catalyst system (e.g. palladium(II) acetate combined with 1,1'-bis(diphenylphosphino)ferrocene) and a reaction-inert solvent (e.g. DMF)

50

9 Compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents OH (which is attached to a CH_2 group) may be prepared by reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $C(O)OR^{7a}$, wherein R^{7a} is as hereinbefore defined, for example at between 0°C and reflux temperature in the presence of a suitable reducing agent (e.g. lithium aluminium hydride) and an appropriate solvent (e.g. THF)

55

10 Compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene,

alkenylene or alkynylene groups are *gem*-disubstituted with two C₁₋₄ alkyl groups (a to D) and are optionally substituted by one or more further substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo or OH, and D represents OH, may be prepared by reaction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C₁₋₃ alkylene, C₂₋₃ alkenylene or C₂₋₃ alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo or OH, and D represents C(O)OR^{7a}, wherein R^{7a} is as hereinbefore defined, with a suitable C₁₋₄ alkyl-delivering organometallic compound (e.g. an alkylmagnesium halide), for example at between -10°C and reflux temperature in the presence of a suitable solvent (e.g. THF)

11 Compounds of formula I in which D represents C(O)N(R⁴)(R⁵), wherein R⁴ and R⁵ are as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula I in which D represents C(O)OR^{7a}, and R^{7a} is as hereinbefore defined, with a compound of formula XI,



or an acid (e.g. HCl) addition salt thereof, wherein R⁴ and R⁵ are as hereinbefore defined, for example at a temperature of between -10 and +150°C and a pressure of between 1 and 10 atmospheres, optionally in the presence (as appropriate) of a Lewis-acidic catalyst (e.g. trimethylaluminium) and a reaction-inert solvent (e.g. toluene)

12 Compounds of formula I in which D represents C(O)N(R⁴)(R⁵), wherein R⁴ and R⁵ are as hereinbefore defined, may alternatively be prepared by reaction of a corresponding compound of formula I in which D represents C(O)OH with a compound of formula XI, as hereinbefore defined, under coupling conditions known to those skilled in the art

13 Compounds of formula I in which D represents C(O)OH may be prepared by hydrolysis of a corresponding compound of formula I in which D represents C(O)OR^{7a}, wherein R^{7a} is as hereinbefore defined, under conditions that are known to those skilled in the art

14 Compounds of formula I in which D represents N(H)R⁶, wherein R⁶ is as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula I in which D represents NH₂ with a compound of formula XII,




wherein R⁶ and L¹ are as hereinbefore defined, for example under conditions that are known to those skilled in the art which include reaction at between -10°C and reflux temperature in the presence of a suitable base (e.g. triethylamine or pyridine) and, optionally, a reaction-inert solvent (e.g. THF or dichloromethane)

15 Compounds of formula I in which A represents C₁₋₄ alkyl and D represents N(R⁴)(R⁵) or N(H)C(O)R^{10a} attached at the 1-, 2- or 3-C (relative to the benzene ring), wherein R⁴, R⁵ and R^{10a} are as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula I in which A represents C₁₋₄ alkenylene unsaturated α,β -, β,γ - or γ,δ - (respectively) relative to the benzene ring and D represents H, with a compound of formula XI, as hereinbefore defined, or a compound of formula XIII,



wherein R^{10a} is as hereinbefore defined, for example at between -10°C and room temperature in the presence of a suitable mercury(II) salt (e.g. mercury(II) acetate, trifluoroacetate, nitrate, or perchlorate), optionally in the presence of a reaction-inert solvent (e.g. THF), and followed by *in situ* reduction of the mercury adduct by the addition of a suitable hydride-delivering agent (e.g. sodium borohydride), optionally in the presence of water.

16 Compounds of formula I in which A represents C₂₋₄ alkylene optionally substituted by one or more substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo or OH, and D represents OH may be prepared by oxidation of a corresponding borane adduct of formula XIV,



XIV

wherein x is 1, 2 or 3 y is (as appropriate) $(3-x)$ or 1. R^{17} is (as appropriate) H, halo, an alkyl, or a cycloalkyl group providing one or two bonds to boron (e.g. diisiamyl or hexyl), A represents (as appropriate) C_{2-4} alkylene optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and R^1 , R^2 , R^3 , X and n are as hereinbefore defined, for example by reaction with a tertiary amine N -oxide (e.g. trimethylamine N -oxide) at between room and reflux temperature in the presence of a reaction-inert solvent (e.g. THF or a THF/diglyme mixture)

The skilled person will appreciate that, in compounds of formula XIV, bonds between boron atoms and piperidine N -atoms may be present

Compounds of formula XIV may be prepared by reaction of a corresponding compound of formula I in which A represents (as appropriate) C_{2-4} alkenylene optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents H with borane or a suitable derivative thereof (e.g. hexylborane, diisiamylborane or 9-borabicyclo[3.3.1]nonane), for example at between -10°C and room temperature in the presence of a suitable solvent (e.g. THF or a THF/diglyme mixture)

17 Compounds of formula I in which A represents a C_{2-4} alkylene group substituted (a to D) with an OH group and D represents OH may be prepared by reaction of a corresponding compound of formula I in which A represents a C_{2-4} alkenylene group and D represents H with a suitable dihydroxylating reagent (e.g. sub-stoichiometric OsO_4 combined with 4-methylmorpholine N -oxide), for example at between 0°C and reflux temperature in the presence of a reaction-inert solvent (e.g. a water/acetone mixture)

18 Compounds of formula I in which A represents a single bond or a C_{1-2} alkylene group (as appropriate) and D represents $\text{C}(\text{O})\text{H}$ may be prepared by reaction of a corresponding compound of formula I in which A represents a C_{2-4} alkylene group substituted (a to D) with an OH group and D represents OH with a reagent that effects 1,2-diol oxidative cleavage (e.g. sodium periodate)

19 Compounds of formula I in which D represents $\text{C}(=\text{NR}^{9a})\text{R}^8$ or $\text{C}(=\text{NOR}^{9b})\text{R}^8$, wherein R^8 , R^{9a} and R^{9b} are as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula I in which D represents $\text{C}(\text{O})\text{R}^8$ with a compound of formula XV.




or a compound of formula XVI



wherein R^{9a} and R^{9b} are as hereinbefore defined, for example under conditions that are known to those skilled in the art which include reaction at between room and reflux temperature in the presence of a suitable solvent (e.g. a lower alkyl alcohol such as methanol or ethanol)

20 Compounds of formula I in which A represents C_{1-4} alkylene substituted (α to D) with an OH group and D

represents $N(H)CH_3$ (at the alkylene chain terminus) may be prepared by reduction of a corresponding compound of formula XVII,




wherein r is 0, 1 or 2, L^2 represents H or a group capable, when attached to a C_2 alkylene unit, of undergoing 1,2-elimination (relative to the L^2 group, e.g. an alkyl or aryl sulfoxide or sulfone), and R^1 , R^2 , R^3 , X and n are as hereinbefore defined, for example, at between -10°C and reflux temperature in the presence of a suitable reducing agent (e.g. lithium aluminium hydride) and a reaction-inert solvent (e.g. THF)

Compounds of formula XVII may be prepared by reaction of a corresponding compound of formula I in which A represents a single bond or C_{1-2} alkylene and D represents $C(O)H$ with a compound of formula XVIII,



wherein L^2 is as hereinbefore defined, for example at between 0°C and reflux temperature in the presence of a suitable solvent (e.g. ethanol) and a catalytic quantity of a cyanide salt (e.g. sodium cyanide)

21 Compounds of formula I wherein R^3 represents C_1 alkyl optionally substituted by C_{3-8} cycloalkyl, Het¹, aryl, adamantyl, (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), or R^3 represents C_{2-10} alkyl, C_{3-10} alkenyl or C_{3-10} alkynyl (which three groups are all optionally substituted by one or more of the relevant substituents identified hereinbefore in respect to R^3), which alkyl, alkenyl or alkynyl groups are attached to the piperidine nitrogen atom via a CH_2 group, wherein Het¹ is as hereinbefore defined, may be prepared by reduction of a corresponding compound of formula XIX,




wherein R^{31} represents H, C_{3-8} cycloalkyl, Het¹, aryl, adamantyl, (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy

and C₁₋₅ alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), C₁₋₉ alkyl, C₂₋₉ alkenyl or C₂₋₉ alkynyl, which alkyl, alkenyl or alkynyl groups are optionally substituted and/or terminated by one or more substituents selected from OR^{11c}, S(O)₂R^{11d}, CN, halo, C₁₋₆ alkoxy carbonyl, C₂₋₄ alkanoyl, C₂₋₆ alkanoyloxy, C₃₋₈ cycloalkyl, C₄₋₉ cycloalkanoyl, N(R^{12a})S(O)₂R¹³, Het¹, aryl, adamantyl (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH₂CN, CONH₂, C₁₋₄ alkyl, C₁₋₄ alkoxy and C₁₋₅ alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), or -W-A¹-N(R^{12b})(R^{12c}), and R¹, R², R^{11c}, R^{11d}, R^{12a} to R^{12c}, R¹³, Het¹, n, p, W, X, A¹, A and D are as hereinbefore defined, using a suitable reducing agent (e.g. lithium aluminium hydride or a borane derivative), for example as described hereinbefore.

The skilled person will appreciate that this reduction may take place simultaneously with other reduction steps described herein (see for example, processes 4, 9 and 16).

Compounds of formula XIX may be prepared by reaction of a corresponding compound of formula XX,



wherein R¹, R², A, D, X and n are as hereinbefore defined with a compound of formula XXI



or a suitable (e.g. carboxylic acid) derivative thereof (e.g. an acid halide or anhydride), wherein R³¹ is as hereinbefore defined using coupling conditions known to those skilled in the art.

Compounds of formulae XIX and XX may be prepared from appropriate precursors by analogy with methods disclosed hereinbefore that describe the preparation of compounds of formula I.

22 Compounds of formula I may be prepared by reaction of a corresponding compound of formula XX, as hereinbefore defined, with a compound of formula VIII, as hereinbefore defined, under conditions that are well known to those skilled in the art, for example as described hereinbefore in respect of the production of compounds of formula VI.

23 Compounds of formula I wherein R³ represents C₁ alkyl, which, in place of being optionally substituted by the substituents as defined hereinbefore, is instead optionally substituted by R³¹, wherein R³¹ is as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula XX, as hereinbefore defined, with a compound of formula XXII



wherein R³¹ is as hereinbefore defined, for example in the presence of a suitable reducing agent (e.g. sodium borohydride, sodium cyanoborohydride or sodium triacetoxyborohydride) and an appropriate solvent (e.g. methanol).

24 Compounds of formula I wherein R³ is a C₁₋₁₀ alkyl, C₄₋₁₀ alkenyl or C₄₋₁₀ alkynyl group that is fully saturated from 1- to 3-C (relative to the piperidine N-atom), and which R³ group is substituted at 2-C (relative to the piperidine N-atom) by S(O)R^{11d}, S(O)₂R^{11d}, alkanoyl, cycloalkanoyl, alkoxy carbonyl, CN, -C(O)-A¹-N(R^{12b})(R^{12c}), -S(O)-A¹-


$N(R^{12b})(R^{12c})$, or $-S(O)_2-A^1-N(R^{12b})(R^{12c})$, wherein R^{11d} , R^{12b} , R^{12c} and A^1 are as hereinbefore defined, may be prepared by reaction of a corresponding compound of formula XX, as hereinbefore defined, with a compound of formula XXIII,



XXIII

wherein R^{3a} represents R^3 as hereinbefore defined except that it does not represent aryl, and that the R^{3a} chain contains an additional carbon-carbon double bond α, β to the Z-substituent, and Z represents $S(O)R^{11d}$, $S(O)_2R^{11d}$, alkanoyl, cycloalkanoyl, alkoxy carbonyl, CN, $-C(O)-A^1-N(R^{12b})(R^{12c})$, $-S(O)-A^1-N(R^{12b})(R^{12c})$, or $-S(O)_2-A^1-N(R^{12b})(R^{12c})$, wherein R^{11d} , R^{12b} , R^{12c} and A^1 are as hereinbefore defined, for example at between room and reflux temperature in the presence of a reaction-inert solvent (e.g. THF)

25. Compounds of formula I in which A represents C_{2-4} alkylene substituted (α to D) with an OH group and D represents $N(R^4)(R^5)$ (at the alkylene chain terminus), and R^4 and R^5 are as hereinbefore defined, may be prepared by reaction of a compound of formula XXIV,




wherein R^1 , R^2 , R^3 , X, n and r are as hereinbefore defined, with a compound of formula XI, as hereinbefore defined, for example at between room and reflux temperature in the presence of a suitable base (e.g. potassium carbonate) and an appropriate solvent (e.g. *N,N*-dimethylformamide)

Compounds of formula XXIV may be prepared by dehydration of a corresponding compound of formula I in which A represents a C_{2-4} alkylene substituted (α to D) with an OH group and D represents OH (at the alkylene chain terminus) under conditions well known to those skilled in the art (e.g. by heating in concentrated sulfuric acid)

Compounds of formula XXIV may alternatively be prepared by epoxidation of a corresponding compound of formula I in which A represents a terminal C_{2-4} alkenylene group and D represents H under conditions well known to those skilled in the art (e.g. by reaction with *meta*-chloroperbenzoic acid)

26. Compounds of formula I in which D represents $N(H)R^4$, wherein R^4 is as hereinbefore defined provided that it does not represent aryl, may be prepared by reduction of a corresponding compound of formula XXV,



wherein R^{4b} and R^{4c} , together with the carbonyl group to which they are attached form a C_{1-6} alkanal, C_{3-6} alkanone, C_{3-8} cycloalkanone, phenyl(C_{1-4})alkanal or phenyl(C_{2-4})alkanone group, which five groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), and R^1 , R^2 , R^3 , A, X and n are as hereinbefore defined (provided that the $-N = C(R^{4b})(R^{4c})$ group is not directly attached to an unsaturated carbon atom), for example at between room and reflux temperature in the presence of a mild reducing agent (e.g. sodium borohydride) and a suitable solvent (e.g. a lower alkyl alcohol such as methanol or ethanol)

Compounds of formula XXV may be prepared by reaction of a corresponding compound of formula I in which D represents NH_2 with a compound of formula XXVI,



wherein R^{4b} and R^{4c} are as hereinbefore defined, for example at between room and reflux temperature in the presence of a reaction-inert solvent (e.g. a lower alkyl alcohol such as methanol or ethanol) and optionally in the presence of a Lewis-acidic catalyst

27. Compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $N(R^4)(R^5)$ (attached to a CH_2 group), wherein R^4 and R^5 are as hereinbefore defined, may be prepared by reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $C(O)N(R^4)(R^5)$, for example in the presence of a suitable reducing agent (e.g. lithium aluminium hydride or a borane derivative) and a reaction-inert solvent (e.g. THF).

Compounds of formulae II, III, VIII, X, R^7aOH , XI, XII, XIII, XV, XVI, XVIII, XXI, XXII, XXIII, XXVI, and derivatives thereof, when not commercially available or not subsequently described, may be obtained either by analogy with the processes described herein or by conventional synthetic procedures, in accordance with standard techniques, from readily available starting materials using appropriate reagents and reaction conditions

[0023] Substituents on alkyl, heterocyclic and aryl groups in the above-mentioned compounds may also be introduced, removed and interconverted using techniques which are well known to those skilled in the art. For example, nitro may be reduced to amino, OH may be alkylated to give alkoxy, alkoxy may be hydrolysed to OH, alkenes may be hydrogenated to alkanes, halo may be hydrogenated to H, etc.

[0024] In some cases it is possible to introduce further substituents into the compounds of formula I directly. For example, chlorination of the phenyl group of compounds of formula I, may be performed by reaction with a solution of chlorine in acetic acid.

[0025] The skilled person will also appreciate that other various standard substituent or functional group interconversions and transformations within certain compounds of formula I will provide other compounds of formula I.

[0026] The compounds of the invention may be isolated from their reaction mixtures using conventional techniques.

[0027] It will be appreciated by those skilled in the art that, in the course of carrying out the processes described above, the functional groups of intermediate compounds may need to be protected by protecting groups.

[0028] Functional groups which it is desirable to protect include oxo, OH, amino and carboxylic acid. Suitable protective groups for oxo include acetals, ketals (e.g. ethylene ketals) and dithianes. Suitable protective groups for OH include trialkylsilyl and diarylalkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl) and tetrahydropyranyl. Suitable protective groups for amino include *tert*-butoxycarbonyl, 9-fluorenylmethoxycarbonyl or benzyloxycarbonyl.

[0029] Suitable protective groups for carboxylic acid include C_{1-6} alkyl or benzyl esters. Suitable protective groups for terminal alkynes include trialkylsilyl and diarylalkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl).

[0030] The protection and deprotection of functional groups may take place before or after any of the reaction steps described hereinbefore.

[0031] Protective groups may be removed in accordance with techniques which are well known to those skilled in the art.

[0032] The use of protecting groups is fully described in "Protective Groups in Organic Chemistry" edited by JWF McOmie, Plenum Press (1973) and "Protective Groups in Organic Synthesis", 2nd edition, TW Greene & PGM Wuts, Wiley-Interscience (1991).

[0033] Persons skilled in the art will also appreciate that, in order to obtain compounds of formula I in an alternative, and, on some occasions, more convenient, manner, the individual process steps mentioned hereinbefore may be performed in a different order, and/or the individual reactions may be performed at a different stage in the overall route (i.e. substituents may be added to and/or chemical transformations performed upon different intermediates to those mentioned hereinbefore in conjunction with a particular reaction). This will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the availability of key intermediates and the protecting group strategy (if any) to be adopted. Clearly, the type of chemistry involved will influence the choice of reagent that is used in the said synthetic steps, the need, and type, of protecting groups that are employed, and the sequence for accomplishing the synthesis. The procedures may be adapted as appropriate to the reactants, reagents and other reaction parameters in a manner that will be evident to the skilled person by reference to standard textbooks and to the examples provided hereinafter.

[0034] It will be appreciated by those skilled in the art that certain protected derivatives of compounds of formula I, which may be made prior to a final deprotection stage, may not possess pharmacological activity as such, but may, in certain instances, be administered orally or parenterally and thereafter metabolised in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". Further, certain compounds of formula I may act as prodrugs of other compounds of formula I.

[0035] It will be further appreciated by those skilled in the art, that certain moieties, known to those skilled in the art as "pro-moieties", for example as described in 'Design of Prodrugs' by H. Bundgaard, Elsevier, 1985 (the disclosure in which document is hereby incorporated by reference), may be placed on appropriate functionalities, when such functionalities are present within compounds of formula I.

[0036] All protected derivatives, and prodrugs, of compounds of formula I are included within the scope of the invention.

[0037] Pharmaceutically acceptable acid addition salts of the compounds of formula I which contain a basic centre may be prepared in a conventional manner. For example, a solution of the free base may be treated with the appropriate acid, either neat or in a suitable solvent, and the resulting salt may then be isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts can be obtained in an analogous manner by treating a solution of a compound of formula I with the appropriate base. Both types of salt may be formed or interconverted using ion-exchange resin techniques.

[0038] The above procedures may be adapted as appropriate to the particular reactants and groups involved and other variants will be evident to the skilled chemist by reference to standard textbooks and to the examples provided hereafter to enable all of the compounds of formula I to be prepared.

[0039] The compounds of the invention are useful because they possess pharmacological activity in animals, especially mammals including humans. They are therefore indicated as pharmaceuticals and, in particular, for use as animal medicaments.

[0040] According to a further aspect of the invention there is provided the compounds of the invention for use as medicaments, such as pharmaceuticals and animal medicaments.

[0041] By the term "treatment", we include both therapeutic (curative) or prophylactic treatment.

[0042] In particular, the compounds of the invention have been found to be useful in the treatment of diseases mediated via opiate receptors, which diseases include irritable bowel syndrome, constipation; nausea; vomiting; pruritus; and conditions characterised by pruritus as a symptom.

[0043] Thus, according to a further aspect of the invention there is provided the use of the compounds of the invention in the manufacture of a medicament for the treatment of a disease mediated via an opiate receptor. There is further provided the use of the compounds of the invention in the manufacture of a medicament for the treatment of irritable bowel syndrome, constipation; nausea, vomiting, pruritus or a medical condition characterised by pruritus as a symptom.

[0044] The compounds of the invention are thus expected to be useful for the curative or prophylactic treatment of pruritic dermatoses including allergic dermatitis and atopy in animals and humans. Other diseases and conditions which may be mentioned include contact dermatitis, psoriasis, eczema and insect bites.

[0045] Thus, the invention provides a method of treating or preventing a disease mediated via an opiate receptor. There is further provided a method of treating irritable bowel syndrome; constipation; nausea; vomiting, pruritus or a medical condition characterised by pruritus as a symptom in an animal (e.g. a mammal), which comprises administering a therapeutically effective amount of a compound of the invention to an animal in need of such treatment.

[0046] The compounds of the invention will normally be administered orally or by any parenteral route, in the form of pharmaceutical preparations comprising the active ingredient, optionally in the form of a non-toxic organic, or inorganic, acid, or base, addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated, as well as the route of administration, the compositions may be administered at varying doses (see below).

[0047] While it is possible to administer a compound of the invention directly without any formulation, the compounds

are preferably employed in the form of a pharmaceutical, or veterinary formulation comprising a pharmaceutically, or veterinarily, acceptable carrier, diluent or excipient and a compound of the invention. The carrier, diluent or excipient may be selected with due regard to the intended route of administration and standard pharmaceutical, and/or veterinary, practice. Pharmaceutical compositions comprising the compounds of the invention may contain from 0.1 percent by weight to 90.0 percent by weight of the active ingredient.

[0048] The methods by which the compounds may be administered for veterinary use include oral administration by capsule, bolus, tablet or drench, topical administration as an ointment, a pour-on, spot-on, dip, spray, mousse, shampoo, collar or powder formulation or, alternatively, they can be administered by injection (e.g. subcutaneously, intramuscularly or intravenously), or as an implant. Such formulations may be prepared in a conventional manner in accordance with standard veterinary practice.

[0049] The formulations will vary with regard to the weight of active compound contained therein, depending on the species of animal to be treated, the severity and type of infection and the body weight of the animal. For parenteral, topical and oral administration, typical dose ranges of the active ingredient are 0.01 to 100 mg per kg of body weight of the animal. Preferably the range is 0.1 to 10 mg per kg.

[0050] The compositions are preferably formulated in a unit dosage form, each dosage containing from about 1 to about 500 mg, more usually about 5 to about 300 mg, of the active ingredient. The term "unit dosage form" refers to physically discreet units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

[0051] In any event, the veterinary practitioner, or the skilled person, will be able to determine the actual dosage which will be most suitable for an individual patient, which may vary with the species, age, weight and response of the particular patient. The above dosages are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

[0052] For veterinary use, the compounds of the invention are of particular value for treating pruritus in domestic animals such as cats and dogs and in horses.

[0053] As an alternative for treating animals, the compounds may be administered with the animal feedstuff and for this purpose a concentrated feed additive or premix may be prepared for mixing with the normal animal feed.

[0054] For human use, the compounds are administered as a pharmaceutical formulation containing the active ingredient together with a pharmaceutically acceptable diluent or carrier. Such compositions include conventional tablet, capsule and ointment preparations which are formulated in accordance with standard pharmaceutical practice.

[0055] Compounds of the invention may be administered either alone or in combination with one or more agents used in the treatment or prophylaxis of disease or in the reduction or suppression of symptoms. Examples of such agents (which are provided by way of illustration and should not be construed as limiting) include antiparasitics, e.g. fipronil, lufenuron, imidacloprid, avermectins (e.g. abamectin, ivermectin, doramectin), milbemycins, organophosphates, pyrethroids, antihistamines, e.g. chlorpheniramine, trimепразине, diphenhydramine, doxylamine, antifungals, e.g. fluconazole, ketoconazole, itraconazole, griseofulvin, amphotericin B, antibacterials, e.g. enrofloxacin, marbofloxacin, ampicillin, amoxycillin, anti-inflammatories, e.g. prednisolone, betamethasone, dexamethasone, carprofen, ketoprofen, dietary supplements, e.g. gamma-linolenic acid and emollients. Therefore, the invention further provides a product containing a compound of the invention and a compound from the above list as a combined preparation for simultaneous, separate or sequential use in the treatment of diseases mediated via opiate receptors.

[0056] The skilled person will also appreciate that compounds of the invention may be taken as a single dose on an "as required" basis (i.e. as needed or desired).

[0057] Thus, according to a further aspect of the invention there is provided a pharmaceutical, or veterinary, formulation including a compound of the invention in admixture with a pharmaceutically, or veterinarily, acceptable adjuvant, diluent or carrier.

[0058] Compounds of the invention may also have the advantage that, in the treatment of human and/or animal patients, they may be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, be more easily absorbed than, or they may have other useful pharmacological properties over, compounds known in the prior art.

[0059] The biological activities of the compounds of the present invention were determined by the following test method.

Biological Test

[0060] Compounds of the present invention have been found to display activity in binding assays selective for the mu opioid receptor in dog brain. The assays were conducted by the following procedure:

[0061] Laboratory bred beagles were used as a source of dog brain tissue. Animals were euthanised, their brains removed and the cerebellum discarded. The remaining brain tissue was sectioned into small pieces approximately 3

g in weight and homogenised in 50 mM Tris pH 7.4 buffer at 4°C using a Kinematica Polytron™ tissue homogeniser. The resulting homogenate was centrifuged at 48,400 x g for 10 minutes and the supernatant discarded. The pellet was resuspended in Tris buffer and incubated at 37°C for 10 minutes. Centrifugation, resuspension and incubation steps were repeated twice more, and the final pellet was resuspended in Tris buffer and stored at -80°C. Membrane material prepared in this manner could be stored for up to four weeks prior to use.

[0062] For mu assays, increasing concentrations of experimental compound, (5×10^{-12} to 10^{-5} M), Tris buffer and ^3H ligand, ([D-Ala², N-Me-Phe⁴, Gly-o⁵]-Enkephalin, DAMGO), were combined in polystyrene tubes. The reaction was initiated by the addition of tissue, and the mixture was incubated at room temperature for 90 minutes. The reaction was terminated by rapid filtration using a Brandel Cell Harvester™ through Betaplate™ GF/A glass fibre filters pre-soaked in 50 mM Tris pH 7.4, 0.1 % polyethylenimine buffer. The filters were then washed three times with 0.5 mL ice-cold Tris pH 7.4 buffer. Washed filters were placed in bags and Starscint™ scintillant added. Bags containing the filters and scintillant were heat sealed and counted by a Betaplate™ 1204 beta counter.

[0063] Duplicate samples were run for each experimental compound and the data generated was analysed using IC₅₀ analysis software in Graphpad Prism. K_i values were calculated using Graphpad Prism according to the following formula

$$K_i = IC_{50} / (1 + [^3\text{H ligand}] / K_D)$$

where IC₅₀ is the concentration at which 50% of the ^3H ligand is displaced by the test compound and K_D is the dissociation constant for the ^3H ligand at the receptor site.

[0064] The invention is illustrated by the following Preparations and Examples in which the following abbreviations may be used

- APCI = atmospheric pressure chemical ionisation
- br (in relation to NMR) = broad
- DMF = N,N-dimethylformamide
- DMSO = dimethylsulfoxide
- d (in relation to time) = day
- d (in relation to NMR) = doublet
- dd (in relation to NMR) = doublet of doublets
- EtOAc = ethyl acetate
- EtOH = ethanol
- h = hour(s)
- m (in relation to NMR) = multiplet
- MeOH = methanol
- min = minute
- q (in relation to NMR) = quartet
- s (in relation to NMR) = singlet
- t (in relation to NMR) = triplet
- THF = tetrahydrofuran

[0065] When reverse phase HPLC is mentioned in the text the following 2 sets of conditions were employed.

[0066] Condition 1 A Phenomenex Magellan™ column, 150 x 21 mm, packed with 5μ C₁₈ silica, eluting with a gradient of acetonitrile 0.1 M aqueous ammonium acetate (30:70 to 95:5 over 10 mins flow rate 20 mL per minute).

[0067] Condition 2 A Dynamax™ column, 42 x 250 mm, packed with 8μ C₁₈ silica, eluting with acetonitrile 0.1 M aqueous ammonium acetate (30:70) at 45 mL per minute.

[0068] In both cases, combination and evaporation of appropriate fractions, determined by analytical HPLC, provided the desired compounds as acetate salts.

[0069] Analytical HPLC conditions used to highlight appropriate fractions were Phenomenex Magellan™ column, 4.6 x 150 mm, packed with 5μ C₁₈ silica, eluting with a gradient of acetonitrile 0.1 M aqueous heptanesulfonic acid (10:90 to 90:10 over 30 min, followed by a further 10 min at 90:10) at 1 mL per minute. Column oven temperature was 40°C, and ultraviolet detection of components was made at 220 nm.

[0070] When column chromatography is referred to this usually refers to a glass column packed with silica gel (40-63 μm). Pressure of ~165 kPa is generally applied and the ratio of crude product : silica gel required for purification is typically 50:1. Alternatively, an Isolute™ SPE (solid phase extraction) column or Waters Sep-Pak™ cartridge packed with silica gel may be used under atmospheric pressure. The ratio of crude product to silica gel required for purification is typically 100:1.

[0071] The hydrochloride salt may be made by methods commonly known to those skilled in the art of synthetic chemistry. Typically, to a solution of free base in dichloromethane (1 g · 100 mL) was added ethereal hydrochloric acid (1.0 M, 1.2 equivalent), the excess solvent was decanted off and the remaining precipitate was washed three times with ether and then dried *in vacuo*.

5 [0072] Melting points were determined using a Gallenkamp melting point apparatus and are uncorrected. ¹H Nuclear magnetic resonance (NMR) spectral data were obtained using a Varian Unity 300 or 400 spectrometer, the observed chemical shifts (δ) being consistent with the proposed structures. Mass spectral (MS) data were obtained on a Fisons Instruments Trio 1000, or a Fisons Instruments Trio 1000 APCI, or a Finnigan Navigator MS, or a Micromass Platform LC spectrometer. The calculated and observed ions quoted refer to the isotopic composition of lowest mass. HPLC means high performance liquid chromatography. Room temperature means 20 to 25°C.

Examples

Example 1: 1-Hexyl-3,4-dimethyl-4-(3-cyanophenyl)piperidine

15 [0073] A solution of 1-hexyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 1, 500 mg, 1.19 mmol) in 1-methyl-2-pyrrolidinone (2.5 mL) was added to a flask containing potassium cyanide (155 mg, 2.38 mmol). The solution was de-oxygenated by evacuating and flushing with nitrogen three times. Catalytic quantities of palladium(II) acetate and 1,1'-bis(diphenylphosphino)ferrocene were added and the reaction mixture was warmed to 20 60°C, at which temperature it was stirred for 3 hours. The reaction mixture was cooled to room temperature and poured into saturated aqueous sodium hydrogencarbonate solution (50 mL). The product was extracted into ethyl acetate (3 x 30 mL). The combined organics were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (15 g) eluted with a gradient of ethyl acetate:hexane 0.880 ammonia (20.79:1 to 50.49:1) to give the title compound as an oil (346 mg).

25 NMR (CDCl₃) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 7.35-7.6 (m, 4H).

MS (electrospray) M/Z (MH⁺) 299.2, C₂₀H₃₀N₂ + H requires 299.2

Example 2: 1-Hexyl-3,4-dimethyl-4-(3-amidophenyl)piperidine

30 [0074] A mixture of polyphosphoric acid (160 mg) and 1-hexyl-3,4-dimethyl-4-(3-cyanophenyl)piperidine (Example 1, 19 mg, 0.064 mmol) was heated at 115°C for one hour. The reaction mixture was then cooled to room temperature and diluted with iced water (0.4 mL). Aqueous sodium hydroxide solution (2 N) was added until the pH was 7. The mixture was extracted with ethyl acetate (3 x 10 mL). The combined organics were dried (Na₂SO₄) and the solvent evaporated *in vacuo* to give a white solid. Purification by column chromatography on silica gel (1 g) eluted with ethyl acetate:ethanol 0.880 ammonia (89.10:1) gave the product as a white solid (6 mg).

35 NMR (CDCl₃, selected data) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.8 (m, 1H), 7.35 (t, 1H), 7.45 (d, 1H), 7.55 (d, 1H), 7.8 (s, 1H).

MS (electrospray) M/Z (MH⁺) 317.3, C₂₀H₃₂N₂O + H requires 317.3

Example 3: 1-Hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine

45 [0075] To a stirred solution of 1-hexyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 1, 500 mg, 1.19 mmol) in anhydrous N,N-dimethylformamide (6 mL) was added triethylamine (1.7 mL, 12.2 mmol) and anhydrous methanol (1.0 mL, 24.7 mmol). The solution was de-oxygenated by evacuating and flushing with nitrogen five times. Palladium(II) acetate (27 mg, 0.12 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (67 mg, 0.12 mmol) were added and the mixture was de-oxygenated again using the same procedure as before. Carbon monoxide gas was bubbled through the mixture for 5 minutes and it was then stirred under an atmosphere of carbon monoxide and heated at 120°C overnight. The solvent was removed *in vacuo* to give a brown oil (0.7 g) which was purified by column chromatography on silica gel (35 g) eluted with a gradient of ethyl acetate:hexane 0.880 ammonia (10.190:1 to 10.90:1 to 25.75:1). This gave the title compound as a yellow oil (250 mg).

50 NMR (CDCl₃) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, containing s, 9H), 1.4-1.55 (m, 2H), 1.7 (m, 1H), 2.1 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.8 (m, 1H), 3.9 (s, 3H), 7.35 (t, 1H), 7.5 (d, 1H), 7.85 (d, 1H), 8.0 (s, 1H).

MS (APCI) M/Z (MH⁺) 332.4, C₂₁H₃₃NO₂ + H requires 332.3

Example 4: 1-Hexyl-3,4-dimethyl-4-(3-(N-isopropyl)amidophenyl)piperidine

55 [0076] In a sealed Wheaton™ vial, 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 40 mg,

0.12 mmol) and isopropylamine (5 mL, 59 mmol) were heated together at 150°C for two days. The reaction mixture was then cooled to room temperature and excess amine was removed by evaporation *in vacuo*. The residue was purified by column chromatography on silica gel (5 g) eluted with ethyl acetate : ethanol : 0.880 ammonia (50:49:1) to give the title compound as an oil (32 mg).

5 NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 15H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 4.3 (m, 1H), 7.3-7.5 (m, 3H), 7.75 (s, 1H).

MS (electrospray) : M/Z (MH⁺) 359.3; C₂₃H₃₈N₂O + H requires 359.3

Example 5: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-butyl)amidophenyl)piperidine

10 [0077] In a sealed Wheaton™ vial, 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 30 mg, 0.09 mmol) and *n*-butylamine (4 mL, 40.5 mmol) were heated together at 140°C for two days. The reaction mixture was then cooled to room temperature and excess amine was removed by evaporation *in vacuo*. The residue was purified by column chromatography on silica gel (5 g) eluted with ethyl acetate : hexane : 0.880 ammonia (40:49:1) to give the title compound as an oil (20 mg).

15 NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 0.95 (t, 3H), 1.2-1.8 (m, 16H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 3.45 (m, 2H), 7.3-7.55 (m, 3H), 7.75 (s, 1H).

MS (electrospray) : M/Z (MH⁺) 373.3; C₂₄H₄₀N₂O + H requires 373.3.

20 Example 6: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-propyl)amidophenyl)piperidine

[0078] In a sealed Wheaton™ vial, 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 30 mg, 0.09 mmol) and *n*-propylamine (4 mL, 94 mmol) were heated together at 140°C for two days. The reaction mixture was then cooled to room temperature and excess amine was removed by evaporation *in vacuo*. The residue was purified by column chromatography on silica gel (5 g) eluted with ethyl acetate : ethanol : 0.880 ammonia (50:49:1) to give the title compound as an oil (3.5 mg).

25 NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.0 (t, 3H), 1.25-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6-1.75 (m, 3H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.45 (m, 2H), 7.3-7.5 (m, 3H), 7.75 (s, 1H).

MS (electrospray) : M/Z (MH⁺) 359.3; C₂₃H₃₈N₂O + H requires 359.3

30 Example 7: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-benzyl)amidophenyl)piperidine

[0079] In a sealed Wheaton™ vial, 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 30 mg, 0.09 mmol) and benzylamine (3 mL, 27.5 mmol) were heated together at 100°C for 76 hours. The reaction mixture was cooled to room temperature, concentrated and the residue purified by column chromatography on silica gel eluted with a gradient of hexane : ethyl acetate (20:80 to 50:50). The title compound was obtained as a pale oil (13 mg).

35 NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (t, 3H), 1.2-1.4 (m, 9H), 1.6-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 4.65 (m, 2H), 7.3-7.4 (m, 6H), 7.45 (d, 1H), 7.55 (d, 1H), 7.8 (s, 1H).

MS (electrospray) : M/Z (MH⁺) 407.3; C₂₇H₃₈N₂O + H requires 407.3

40 Example 8: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine

[0080] In a sealed Wheaton™ vial, 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 30 mg, 0.09 mmol) and ethylamine (3 mL, 45.8 mmol) were heated together at 100°C for 60 hours. The ethylamine was found to be evaporating, thus the reaction mixture was transferred to a sealed bomb and heated at 100°C and 690 kPa for a further 16 hours. The reaction mixture was cooled to room temperature, concentrated and then purified by column chromatography on silica gel eluted with a gradient of hexane : ethyl acetate (20:80 to 50:50). The title compound was obtained as a pale oil (11 mg).

45 NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 12H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 3.5 (m, 2H), 7.35 (t, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H).

50 MS (thermospray) : M/Z (MH⁺) 345.1; C₂₂H₃₆N₂O + H requires 345.3.

Example 9: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-isobutyl)amidophenyl)piperidine

55 [0081] The title compound was prepared by the method of Example 7, substituting benzylamine with isobutylamine (3 mL, 30.18 mmol). This gave the title compound as a pale oil (9 mg).

NMR (CDCl₃, selected data) : 0.75 (d, 3H), 0.9 (t, 3H), 1.0 (d, 6H), 1.2-1.35 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 1.9 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.3 (t, 2H), 7.35 (t, 1H), 7.45 (d, 1H), 7.5

(d, 1H), 7.75 (s, 1H)

MS (electrospray) M/Z (MH^+) 373.3, $C_{24}H_{40}N_2O + H$ requires 373.3

Example 10: 1-Hexyl-3,4-dimethyl-4-(3-[*N*-(2-methoxyethyl)]amidophenyl)piperidine

[0082] In a sealed Wheaton™ vial 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 30 mg, 0.09 mmol) and 2-methoxyethylamine (3 mL, 34.5 mmol) were heated together at 130°C for 120 hours. The reaction mixture was cooled to room temperature, concentrated and purified by column chromatography on silica gel eluted with a gradient of hexane:ethyl acetate:0.680 ammonia gradient (10:89:1 to 49:50:1). This gave the title compound as a pale oil (8 mg).

NMR ($CDCl_3$, selected data) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.3-3.7 (m, 7H), 7.35 (t, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H).
MS (APCI) M/Z (MH^+) 374.5, $C_{23}H_{38}N_2O_2 + H$ requires 374.3

Example 11: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-methyl)amidophenyl)piperidine

[0083] To a suspension of anhydrous methylamine hydrochloride (17 mg, 0.25 mmol) in anhydrous toluene (0.5 mL) stirred under nitrogen and cooled in an ice bath was added a solution of trimethylaluminum (2.0 M in toluene, 0.12 mL, 0.24 mmol). The mixture was allowed to warm to room temperature while stirring for 4 hours, then it was treated with a solution of 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 40 mg, 0.12 mmol) in anhydrous toluene (1.5 mL). The resulting mixture was heated at reflux overnight, then quenched with dilute hydrochloric acid (10 mL of 2 *N*) and extracted with diethyl ether (10 mL). The aqueous phase was basified to pH 13 with aqueous sodium hydroxide solution (2 *N*) and extracted with dichloromethane (3 x 20 mL). The combined dichloromethane extracts were dried (Na_2SO_4) and concentrated *in vacuo* to give an orange oil (31 mg) which was purified by column chromatography on silica gel (1.2 g) eluted with ethyl acetate:hexane:0.880 ammonia (50:50:1). This gave the title compound as a colourless residue (17 mg).

NMR ($CDCl_3$, selected data) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.0 (d, 3H), 7.35 (t, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H).
MS (APCI) M/Z (MH^+) 331.1, $C_{21}H_{34}N_2O + H$ requires 331.3

Example 12: 1-Hexyl-3,4-dimethyl-4-(3-(*N,N*-dimethyl)amidophenyl)piperidine

[0084] The title compound was prepared as for Example 11 except using anhydrous dimethylamine hydrochloride (20 mg, 0.25 mmol) in place of methylamine hydrochloride. This gave a pale yellow oil (38 mg).

NMR ($CDCl_3$) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 2.95 (br s, 3H), 3.1 (br s, 3H), 7.15-7.25 (m, 1H), 7.25-7.4 (m, 3H).
MS (thermospray) M/Z (MH^+) 345.2, $C_{22}H_{36}N_2O + H$ requires 345.3

Example 13: 1-Hexyl-3,4-dimethyl-4-(3-(*N,N*-diethyl)amidophenyl)piperidine

[0085] To a suspension of anhydrous diethylamine hydrochloride (30 mg, 0.27 mmol) in anhydrous toluene (0.5 mL) stirred under nitrogen and cooled in an ice bath was added a solution of trimethylaluminum (2.0 M in toluene, 0.14 mL, 0.28 mmol). The mixture was allowed to warm to room temperature while stirring for 2 hours, then it was treated with a solution of 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 44 mg, 0.13 mmol) in anhydrous toluene (1 mL). The resulting mixture was heated at reflux for 2½ days, then quenched with dilute hydrochloric acid (10 mL of 2 *N*) and extracted with diethyl ether (10 mL). The aqueous phase was basified to pH 13 with aqueous sodium hydroxide solution (2 *N*) and extracted with dichloromethane (3 x 20 mL). The combined dichloromethane extracts were dried (Na_2SO_4) and concentrated *in vacuo* to give an orange oil (52 mg) which was purified by column chromatography on silica gel (1.5 g) eluted with a gradient of ethyl acetate:hexane:0.880 ammonia (10:90:1 to 20:40:1). This gave the title compound as a yellow oil (34 mg).

NMR ($CDCl_3$) 0.75 (d, 3H), 0.9 (m, 3H), 1.0-1.4 (m, 15H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.25 (br m, 2H), 3.55 (br m, 2H), 7.1-7.2 (m, 1H), 7.2-7.35 (m, 3H).
MS (APCI) M/Z (MH^+) 373.1, $C_{24}H_{40}N_2O + H$ requires 373.3

Example 14: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-*tert*-butyl)amidophenyl)piperidine

[0086] A solution of 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 40 mg, 0.12 mmol) in dilute hydrochloric acid (5 mL of 2 *N*) was heated at reflux overnight. The solvent was removed *in vacuo* and the residue

was taken up in methanol and re-concentrated *in vacuo* to give a brown oil (40 mg) which was dissolved in dichloromethane (1 mL) and treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (25 mg, 0.13 mmol), *N*-methylmorpholine (27 μ L, 0.25 mmol) and *tert*-butylamine (14 μ L, 0.13 mmol). The resulting mixture was stirred at room temperature overnight, then poured into saturated aqueous sodium hydrogencarbonate solution (5 mL) and extracted with dichloromethane (3 x 5 mL). The combined extracts were dried (Na_2SO_4) and concentrated *in vacuo* to give a brown residue which was purified by column chromatography on silica gel (2.5 g) eluted with a gradient of ethyl acetate : hexane : 0.880 ammonia (5:95:1 to 10:90:1 to 20:80:1 to 30:70:1). This gave the title compound as a colourless residue (10 mg).

NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 11H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.8 (m, 1H), 7.3-7.45 (m, 3H), 7.75 (s, 1H).
MS (APCI) : M/Z (MH^+) 373.1; $\text{C}_{24}\text{H}_{40}\text{N}_2\text{O} + \text{H}$ requires 373.3.

Example 15: 1-Benzyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine

[0087] A stirred solution of 1-benzyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 2, 506 mg, 1.18 mmol), triethylamine (1.6 mL, 11.8 mmol) and anhydrous methanol (1.9 mL, 46.8 mmol) in anhydrous *N,N*-dimethylformamide (6 mL) was de-oxygenated by evacuating and flushing with nitrogen five times. Palladium(II) acetate (30 mg, 0.13 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (63 mg, 0.11 mmol) were added and the mixture was again de-oxygenated, using the same method as before. Carbon monoxide gas was bubbled through the mixture for ca 5 minutes and it was subsequently heated at 80°C under an atmosphere of carbon monoxide overnight. The mixture was then poured into water (100 mL) and extracted with diethyl ether (3 x 100 mL). The combined extracts were dried (Na_2SO_4) and concentrated *in vacuo* to give an orange oil (250 mg). A black residue remaining in the reaction flask, insoluble in diethyl ether, was dissolved in dichloromethane and transferred to the separating funnel containing the aqueous layer and this was re-extracted with dichloromethane (3 x 50 mL). The combined organics were filtered through Celite® to remove residual palladium, dried (Na_2SO_4) and concentrated *in vacuo* to give a brown oil (270 mg). The combined oils were purified by silica (25 g) column chromatography eluting with a gradient of hexane : ethyl acetate : 0.880 ammonia (140:10:1 to 90:10:1) to give the title compound as a colourless oil (335 mg).
NMR (CDCl_3) : 0.75 (d, 3H), 1.35 (s, 3H), 1.65 (m, 1H), 2.05 (m, 1H), 2.3-2.5 (m, 2H), 2.5-2.65 (m, 2H), 2.85 (m, 1H), 3.45 (d, 1H), 3.6 (d, 1H), 3.9 (s, 3H), 7.2-7.4 (m, 6H), 7.5 (d, 1H), 7.85 (d, 1H), 8.0 (s, 1H).
MS (thermospray) : M/Z (MH^+) 338.2; $\text{C}_{22}\text{H}_{27}\text{NO}_2 + \text{H}$ requires 338.2.

Example 16: 1-Benzyl-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine

[0088] A stirred suspension of ethylamine hydrochloride (880 mg, 10.8 mmol) in anhydrous toluene (10 mL) was de-oxygenated by evacuating and flushing with nitrogen three times. Stirring under nitrogen it was cooled in an ice bath and treated with trimethylaluminium solution (2.0 M in toluene, 5.4 mL, 10.8 mmol) via syringe. The mixture was allowed to warm to room temperature while stirring for 1½ hours. A solution of 1-benzyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 15, 1.81 g, 5.36 mmol) in anhydrous toluene (20 mL) was added via syringe and the reaction mixture was heated at reflux overnight. After it had cooled, aqueous hydrochloric acid (100 mL of 2 M) was added and the mixture was extracted with diethyl ether (100 mL). The organic extract was back-washed with aqueous hydrochloric acid (50 mL of 2 M). The combined aqueous phases were basified to pH 13 with aqueous sodium hydroxide solution (2 M) and then extracted with dichloromethane (300 mL followed by 2 x 100 mL). The combined dichloromethane extracts were washed with water (150 mL) followed by saturated aqueous sodium chloride solution (150 mL), dried (Na_2SO_4) and concentrated *in vacuo* to give a brown oil (2.0 g). Purification by silica (100 g) column chromatography eluting with a gradient of ethyl acetate : hexane : 0.880 ammonia (20:80:1 to 30:70:1 to 40:60:1) gave the title compound as a cream foam (805 mg).
NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 1.25 (t, 3H), 1.35 (s, 3H), 1.65 (m, 1H), 2.05 (m, 1H), 2.35-2.45 (m, 2H), 2.5-2.6 (m, 2H), 2.85 (m, 1H), 3.4-3.55 (m, 3H), 3.6 (d, 1H), 7.2-7.35 (m, 6H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H).
MS (APCI) : M/Z (MH^+) 351.3; $\text{C}_{23}\text{H}_{30}\text{N}_2\text{O} + \text{H}$ requires 351.2.

Example 17: 1-(2-Phenoxyethyl)-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine

[0089] A stirred mixture of 3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine (Preparation 3, 50 mg, 0.19 mmol), 2-bromoethyl phenyl ether (42 mg, 0.20 mmol), and sodium hydrogencarbonate (19.1 mg, 0.22 mmol) in anhydrous *N,N*-dimethylformamide (1 mL) was heated at 100°C for 2 hours. The solvent was then removed *in vacuo* to give a brown oil which was purified by column chromatography on silica gel (4 g) eluted initially with CH_2Cl_2 , changing incrementally to CH_2Cl_2 : MeOH (25:1), to give the title compound as a yellow oil (55 mg).
NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 1.25 (t, 3H), 1.35 (s, 3H), 1.65 (m, 1H), 2.05 (m, 1H), 2.35 (m, 1H), 2.55

(m, 1H), 2.6-2.8 (m, 3H), 2.8-2.95 (m, 2H), 3.5 (m, 2H), 4.1 (m, 2H), 6.9-7.0 (m, 3H), 7.3 (m, 2H), 7.35 (m, 1H), 7.4 (m, 1H), 7.5 (d, 1H), 7.75 (s, 1H)
 MS (thermospray) M/Z (MH^+) 381.2, $C_{24}H_{32}N_2O_2 + H$ requires 381.3

5 **Example 18: 1-(5-Methylhexyl)-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine**

[0090] A stirred mixture of 3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine (Preparation 3, 50 mg, 0.19 mmol), 1-bromo-5-methylhexane (37 mg, 0.20 mmol), and sodium hydrogencarbonate (19.1 mg, 0.22 mmol) in anhydrous *N,N*-dimethylformamide (1 mL) was heated at 100°C for 2½ hours. The solvent was then removed *in vacuo* to give a brown oil which was purified by column chromatography on silica gel (4 g) eluted initially with CH_2Cl_2 , changing incrementally to CH_2Cl_2 / MeOH (25/1) to give the title compound as a yellow oil (58 mg).
 10 NMR ($CDCl_3$, selected data) 0.75 (d, 3H), 0.85 (d, 6H), 1.15 (m, 2H), 1.2-1.35 (m, 8H), 1.35-1.6 (m, 3H), 1.65 (m, 1H), 2.05 (s, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 3.5 (m, 2H), 7.35 (t, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H)
 15 MS (thermospray) M/Z (MH^+) 359.5, $C_{23}H_{38}N_2O + H$ requires 359.3

Example 19: 1-(3-Phenylpropyl)-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine

[0091] A stirred mixture of 3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine (Preparation 3, 50 mg, 0.19 mmol), 1-bromo-3-phenylpropane (32 µL, 0.21 mmol), and sodium hydrogencarbonate (19.1 mg, 0.22 mmol) in anhydrous *N,N*-dimethylformamide (1 mL) was heated at 100°C for 2 hours. The solvent was then removed *in vacuo* to give a brown oil which was purified by column chromatography on silica gel (4 g) eluted initially with CH_2Cl_2 , changing incrementally to CH_2Cl_2 / MeOH (25/1), followed by further purification by column chromatography on silica gel (4 g) eluted with a gradient of CH_2Cl_2 / MeOH (50/1 to 50/2). This gave the title compound as a yellow oil (55 mg).
 20 NMR ($CDCl_3$, selected data) 0.75 (d, 3H), 1.25 (t, 3H), 1.3 (s, 3H), 1.65 (m, 1H), 1.8 (m, 2H), 2.05 (m, 1H), 2.25-2.45 (m, 4H), 2.45-2.7 (m, 4H), 2.85 (m, 1H), 3.5 (m, 2H), 7.15-7.2 (m, 3H), 7.25 (m, 2H), 7.35 (t, 1H), 7.4 (d, 1H), 7.5 (d, 1H), 7.75 (s, 1H)
 MS (thermospray) M/Z (MH^+) 379.0, $C_{25}H_{34}N_2O + H$ requires 379.3

30 **Example 20: 1-(5-Cyanopentyl)-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine**

[0092] A stirred mixture of 3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine (Preparation 3, 50 mg, 0.19 mmol), 6-bromohexanenitrile (28 µL, 0.20 mmol) and sodium hydrogencarbonate (19.1 mg, 0.22 mmol) in anhydrous *N,N*-dimethylformamide (1 mL) was heated at 100°C for 4 hours. The solvent was then removed *in vacuo* to give a brown oil which was purified by column chromatography on silica gel (4 g) eluted initially with CH_2Cl_2 , changing incrementally to CH_2Cl_2 / MeOH (25/1), to give a yellow oil (82 mg), followed by further purification by column chromatography on silica gel (4 g) eluted with a gradient of CH_2Cl_2 / MeOH (50/1 to 50/2). This gave the title compound as a yellow oil (54 mg).
 35 NMR ($CDCl_3$, selected data) 0.8 (d, 3H), 1.25 (t, 3H), 1.35 (s, 3H), 1.5 (m, 2H), 1.55-1.8 (m, 5H), 2.15 (m, 1H), 2.35 (t, 2H), 2.4-2.6 (m, 4H), 2.6-2.75 (m, 2H), 2.95 (m, 1H), 3.5 (m, 2H), 7.3-7.45 (m, 2H), 7.5 (d, 1H), 7.75 (s, 1H)
 40 MS (thermospray) M/Z (MH^+) 356.4, $C_{22}H_{33}N_3O + H$ requires 356.3

Example 21: 1-Hexyl-3,4-dimethyl-4-(3-aminomethylphenyl)piperidine

[0093] To a stirred solution of 1-hexyl-3,4-dimethyl-4-(3-cyanophenyl)piperidine (Example 1, 800 mg, 2.68 mmol) in anhydrous tetrahydrofuran (40 mL) at 0°C under nitrogen was added lithium aluminium hydride (1.0 M in tetrahydrofuran, 4.0 mL, 4.0 mmol). The reaction mixture was allowed to warm to room temperature, before being heated to 37°C for 30 minutes. Subsequently, diethyl ether (50 mL) then aqueous sodium hydroxide (0.3 mL, 15% w/v solution) and finally water (0.45 mL) were added. The white solid formed was filtered off. The filtrate was washed with saturated aqueous sodium hydrogencarbonate solution (2 x 50 mL). The aqueous phases were back-extracted with diethyl ether (50 mL). The combined organics were dried ($MgSO_4$) and concentrated *in vacuo* to give the title compound as an oil (770 mg).
 45 NMR ($CDCl_3$, selected data) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.8 (m, 1H), 3.85 (s, 2H), 7.1-7.35 (m, 4H)
 50 MS (APCI) M/Z (MH^+) 303.4, $C_{20}H_{34}N_2 + H$ requires 303.3

Example 22: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-methoxycarbonyl)aminomethylphenyl)piperidine

- [0094] To a stirred solution of 1-hexyl-3,4-dimethyl-4-(3-aminomethylphenyl)piperidine (Example 21, 100 mg, 0.33 mmol) in pyridine (2 mL, dried over basic alumina) at 0°C under nitrogen was added methyl chloroformate (40 μ L, 0.52 mmol). The reaction mixture was stirred at room temperature under a nitrogen atmosphere overnight. Subsequently, ice was added followed by aqueous sodium hydroxide (5 *N* solution) to give a pH of 11. The mixture was extracted with diethyl ether (3 x 25 mL). The combined organic phases were dried (MgSO_4) and then concentrated *in vacuo* at 70°C to give an oil (104 mg) which was purified by column chromatography on silica gel (29 g) eluted with a gradient of ethyl acetate : hexane (1.2 to 2 : 1). This gave the title compound as an oil (65 mg).
- 10 NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.35 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.4 (m, 4H), 2.4-2.6 (m, 2H), 2.8 (m, 1H), 3.7 (s, 3H), 4.35 (d, 2H), 7.1 (d, 1H), 7.15-7.35 (m, 3H).
- MS (thermospray) : M/Z (MH^+) 361.4; $\text{C}_{22}\text{H}_{36}\text{N}_2\text{O}_2 + \text{H}$ requires 361.3

Example 23: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-acetyl)aminomethylphenyl)piperidine

- 15 [0095] This preparation was carried out using the procedure described for Example 22 except using acetyl chloride (35 μ L, 0.49 mmol) in place of methyl chloroformate. This gave the title compound as an oil (100 mg).
- NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 1.95-2.05 (m, 4H), 2.2-2.4 (m, 4H), 2.4-2.6 (m, 2H), 2.8 (m, 1H), 4.4 (d, 2H), 7.1 (d, 1H), 7.15-7.35 (m, 3H).
- 20 MS (APCI) : M/Z (MH^+) 345.3; $\text{C}_{22}\text{H}_{36}\text{N}_2\text{O} + \text{H}$ requires 345.3

Example 24: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-methanesulfonyl)aminomethylphenyl)piperidine

- 25 [0096] This preparation was carried out using the procedure described for Example 22 except using methanesulfonyl chloride (40 μ L, 0.52 mmol) in place of methyl chloroformate. This gave the title compound as an oil (94 mg).
- NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.4 (m, 4H), 2.4-2.65 (m, 2H), 2.75-2.9 (m, 4H), 4.3 (s, 2H), 7.15 (d, 1H), 7.2-7.35 (m, 3H).
- MS (APCI) : M/Z (MH^+) 381.6; $\text{C}_{21}\text{H}_{36}\text{N}_2\text{O}_2\text{S} + \text{H}$ requires 381.3.

Example 25: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-trifluoromethanesulfonyl)aminomethylphenyl)piperidine

- 30 [0097] To a solution of 1-hexyl-3,4-dimethyl-4-(3-aminomethylphenyl)piperidine (Example 21, 100 mg, 0.33 mmol) in pyridine (1.5 mL, dried over basic alumina) stirred under nitrogen was added trifluoromethanesulfonyl chloride (0.3 mL, 0.28 mmol). The reaction mixture was stirred at room temperature under a nitrogen atmosphere overnight. Aqueous sodium hydroxide (20 mL of 2 *N*) was added and the mixture was extracted with dichloromethane (3 x 20 mL). The aqueous layer was treated with dilute aqueous hydrochloric acid (20 mL of 1 *N*) and then extracted with dichloromethane (2 x 20 mL). The organic phases were combined, dried (MgSO_4) and concentrated *in vacuo* to give an orange oil (40 mg) which was purified by column chromatography on silica gel (1.2 g) eluted with a gradient of ethyl acetate : hexane : triethylamine (20:80:1) to ethyl acetate : triethylamine (100:1). This gave the title compound as an oil (30 mg).
- 40 NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.75 (m, 6H), 2.8 (m, 1H), 4.4 (s, 2H), 7.15 (d, 1H), 7.2-7.35 (m, 3H).
- MS (thermospray) : M/Z (MH^+) 435.1; $\text{C}_{21}\text{H}_{33}\text{F}_3\text{N}_2\text{O}_2\text{S} + \text{H}$ requires 435.2.

Example 26: 1-Hexyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine

- 45 [0098] A solution of 1-hexyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 1, 1.5 g, 3.6 mmol) in 1,4-dioxane (17 mL) was de-oxygenated by evacuating and flushing with nitrogen five times. Vinyl tributyltin (1.06 mL, 3.71 mmol) was added under stirring, followed by lithium chloride (456 mg, 10.76 mmol), tetrakis(triphenylphosphine)palladium(0) (catalytic) and 2,6-di-*tert*-butyl-4-methylphenol (2 crystals). The suspension was stirred under nitrogen and heated at reflux for ten hours. After cooling to room temperature the reaction mixture was quenched with aqueous ammonium hydroxide solution (50 mL, 1.0 *M*) and further diluted with ethyl acetate (50 mL). The phases were separated and the aqueous layer was further extracted with ethyl acetate (3 x 25 mL). The combined organics were dried (Na_2SO_4) and concentrated *in vacuo*. The residual oil was purified by column chromatography on silica gel (120 g) eluted with ethyl acetate : hexane : ammonia (39:60:1) to give the title compound as an oil (890 mg).
- 50 NMR (CDCl_3) : 0.75 (d, 3H), 0.85 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 5.25 (d, 1H), 5.75 (d, 1H), 6.7 (dd, 1H), 7.1-7.35 (m, 4H).
- MS (thermospray) : M/Z (MH^+) 300.4; $\text{C}_{21}\text{H}_{33}\text{N} + \text{H}$ requires 300.3.

Example 27: 1-Hexyl-3,4-dimethyl-4-(3-(1,2-dihydroxyethyl)phenyl)piperidine

[0099] 1-Hexyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine (Example 26, 200 mg, 0.67 mmol) was dissolved in a mixture of water (2 mL) and acetone (18 mL). 4-methylmorpholine N-oxide (172 mg, 1.47 mmol) was added with stirring followed by osmium tetroxide (200 μ L, 2.5% w/w in *tert*-butanol). The reaction mixture was stirred at room temperature for 4 hours before the solvent was removed by evaporation *in vacuo*. The residue was partitioned between dichloromethane (25 mL) and water (25 mL). The organic phase was separated and dried (Na_2SO_4). Concentration *in vacuo* gave a residue which was purified by column chromatography on silica gel (10 g) eluted with a gradient of ethyl acetate/hexane/ammmonium hydroxide solution (50/49/1 to 60/33/1), followed by ethyl acetate/methanol/ammmonium hydroxide solution (94/5/1). Combination of the appropriate fractions and evaporation to dryness *in vacuo* gave the product as a yellow oil (145 mg).

NMR (CDCl_3 , selected data): 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 3.7 (m, 2H), 4.80 (m, 1H), 7.1-7.4 (m, 4H).

MS (thermospray) M/Z (MH^+) 334.5, $\text{C}_{21}\text{H}_{35}\text{NO}_2 + \text{H}$ requires 334.3

Example 28: 1-Hexyl-3,4-dimethyl-4-(3-formylphenyl)piperidine

[0100] 1-Hexyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine (Example 26, 200 mg, 0.67 mmol) was dissolved in a mixture of water (2 mL) and acetone (18 mL). Osmium tetroxide (200 μ L, 2.5% w/w in *tert*-butanol) was added followed by sodium periodate (572 mg, 2.68 mmol) which was added portionwise. The reaction mixture was stirred at room temperature for 26 hours, then it was filtered to remove precipitate and the solvent was removed by evaporation *in vacuo*. The residue was partitioned between dichloromethane (25 mL) and saturated sodium chloride solution (25 mL). The organic phase was separated, dried (Na_2SO_4) and the solvent removed *in vacuo*. The residue was purified by column chromatography on silica gel (50 g) eluted with ethyl acetate/hexane/0.880 ammonia (74/25/1). The title compound was obtained as an oil (80 mg).

NMR (CDCl_3 , selected data): 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.6 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.85 (m, 1H), 7.5 (t, 1H), 7.55 (d, 1H), 7.7 (d, 1H), 7.8 (s, 1H), 10.0 (s, 1H).

MS (electrospray) M/Z (MH^+) 302.0, $\text{C}_{20}\text{H}_{31}\text{NO} + \text{H}$ requires 302.2

Example 29: 1-Hexyl-3,4-dimethyl-4-(3-(*N*-hydroxy)iminomethylphenyl)piperidine

[0101] A solution of 1-hexyl-3,4-dimethyl-4-(3-formylphenyl)piperidine (Example 28, 80 mg, 0.27 mmol) in a mixture of pyridine (1 mL) and ethanol (1 mL) was treated with hydroxylamine hydrochloride (22 mg, 0.32 mmol) and the resulting mixture was heated at reflux for 18 hours. The solvent was evaporated *in vacuo* and the residual orange oil was purified by column chromatography on silica gel (10 g) eluted with a gradient of dichloromethane/methanol/0.380 ammonia (98/1/1 to 94/5/1). This gave the title compound as an oil (18 mg).

NMR (CDCl_3 , selected data): 0.6 (d, 3H), 0.85 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.6 (m, 2H), 1.7 (m, 1H), 2.1 (m, 1H), 2.2-2.75 (m, 6H), 2.95 (m, 1H), 7.2-7.4 (m, 3H), 7.6 (s, 1H), 8.1 (s, 1H).

MS (thermospray) M/Z (MH^+) 317.6, $\text{C}_{20}\text{H}_{32}\text{N}_2\text{O} + \text{H}$ requires 317.3

Example 30: 1-Hexyl-3,4-dimethyl-4-(3-acetylphenyl)piperidine

[0102] To a solution of 1-hexyl-3,4-dimethyl-4-(3-cyanophenyl)piperidine (Example 1, 791 mg, 2.65 mmol) in anhydrous tetrahydrofuran (6 mL) at 0°C was added methyl lithium (2.46 mL, 3.45 mmol) and the mixture darkened. The solution was then warmed to room temperature and stirred under a nitrogen atmosphere for 1 hour before being poured onto water (10 mL). The basic aqueous layer was extracted with diethyl ether/ethyl acetate (1/1, 3 x 10 mL). The combined organic extracts were dried (MgSO_4) and concentrated *in vacuo*. This gave the crude title compound as a colourless oil (720 mg, 86%).

NMR (CDCl_3 , selected data): 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.6 (s, 3H), 2.85 (m, 1H), 7.4 (t, 1H), 7.5 (d, 1H), 7.75 (d, 1H), 7.95 (s, 1H).

MS (thermospray) M/Z (MH^+) 316.3, $\text{C}_{21}\text{H}_{33}\text{NO} + \text{H}$ requires 316.3

Example 31: 1-Hexyl-3,4-dimethyl-4-(3-ethynylphenyl)piperidine

[0103] A solution of 1-hexyl-3,4-dimethyl-4-(3-[2-(trimethylsilyl)ethynyl]phenyl)piperidine (Preparation 4, 150 mg, 0.40 mmol) in tetrahydrofuran (2 mL) was cooled to -70°C and tetrabutylammonium fluoride (1.0 M in THF, 0.41 mL, 0.41 mmol) was added slowly. The reaction mixture was allowed to warm to room temperature gradually before being diluted with dichloromethane (10 mL) and water (10 mL). The phases were separated and the aqueous layer was

further extracted with dichloromethane (2 x 10 mL). The combined organics were dried (Na_2SO_4) and the solvent evaporated *in vacuo*. The oily yellow residue was purified by column chromatography on silica gel (10 g) eluted with ethyl acetate : hexane : 0.880 ammonia (10:89:1) to give the title compound as an oil (100 mg).

NMR (CDCl_3) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.35 (m, 9H), 1.35-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.6 (m, 6H), 2.8 (m, 1H), 3.05 (s, 1H), 7.2-7.35 (m, 3H), 7.45 (s, 1H).

MS (APCI) : M/Z (MH^+) 298.6; $\text{C}_{21}\text{H}_{31}\text{N} + \text{H}$ requires 298.3

Example 32: 1-Hexyl-3,4-dimethyl-4-(3-(1,1-dimethyl)hydroxymethylphenyl)piperidine

[0104] A solution of 1-hexyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Example 3, 50 mg, 0.15 mmol) in anhydrous tetrahydrofuran was de-oxygenated by evacuating and flushing with nitrogen three times. The solution was then cooled to 0°C and treated with methylmagnesium chloride (0.5 mL, 1.5 mmol, 3.0 M in tetrahydrofuran) dropwise. The reaction mixture was stirred at 50°C for 2 hours and then saturated aqueous ammonium chloride solution (20 mL) was added followed by saturated aqueous sodium hydrogencarbonate (20 mL). The mixture was extracted with ethyl acetate (3 x 15 mL) and the combined organic phases were dried (MgSO_4) and concentrated *in vacuo* to give an oil (39 mg). The residue was purified by column chromatography on silica gel (1 g) eluted with a gradient of ethyl acetate : hexane : ammonia (50:50:1 to 25:75:1) to give the title compound as a colourless oil (30 mg).

NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.25-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (s, 6H), 1.65 (m, 1H), 2.05 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 7.15 (m, 1H), 7.2-7.3 (m, 2H), 7.45 (s, 1H).

MS (APCI) : M/Z (MH^+) 332.4; $\text{C}_{22}\text{H}_{37}\text{NO} + \text{H}$ requires 332.3

Example 33: 1-Hexyl-3,4-dimethyl-4-(3-hydroxymethylphenyl)piperidine

[0105] A stirred solution of 1-hexanoyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine (Preparation 7, 80 mg, 0.23 mmol) in anhydrous tetrahydrofuran (1 mL) under nitrogen was treated with lithium aluminium hydride (1.0 M in ether, 0.70 mL, 0.70 mmol) and the mixture was stirred at room temperature for 3 hours. The reaction mixture was then quenched with water (7.5 mL) and extracted with ethyl acetate (7 mL). The phases were separated and the aqueous layer was further extracted with ethyl acetate (2 x 5 mL). The combined organics were dried (Na_2SO_4) and the solvent removed *in vacuo* to give the title compound as a pale oil (30 mg).

NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.65 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 4.7 (s, 2H), 7.1-7.35 (m, 4H).

MS (thermospray) : M/Z (MH^+) 304.3; $\text{C}_{20}\text{H}_{33}\text{NO} + \text{H}$ requires 304.3.

Example 34: 1-Hexyl-3,4-dimethyl-4-(3-(2-hydroxyethyl)phenyl)piperidine

[0106] To a stirred solution of 1-hexanoyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine (Preparation 8, 50 mg, 0.16 mmol) in bis(2-methoxyethyl)ether (1.5 mL) at 0°C under nitrogen was added dropwise borane (1.0 M in tetrahydrofuran, 0.35 mL, 0.35 mmol). The reaction mixture was stirred at 0°C for 30 minutes and then for 2 hours at room temperature. Trimethylamine N-oxide (48 mg, 0.64 mmol) was subsequently added and the reaction mixture heated at reflux under a nitrogen atmosphere for 2 hours. To the cooled reaction was then added diethyl ether (10 mL) and saturated aqueous sodium chloride solution (10 mL). The phases were separated and the aqueous layer was further extracted with diethyl ether (10 mL). The combined organics were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (1.5 g) eluted with ethyl acetate : hexane (50:50) to give the title compound as an oil (30 mg).

NMR (CDCl_3 , selected data) : 0.75 (d, 3H), 0.9 (m, 3H), 1.25-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.6 (m, 2H), 2.8 (m, 1H), 2.85 (t, 2H), 3.85 (t, 2H), 7.05 (d, 1H), 7.1-7.35 (m, 3H).

MS (APCI) : M/Z (MH^+) 318.6; $\text{C}_{21}\text{H}_{35}\text{NO} + \text{H}$ requires 318.3.

Example 35: 1-Hexyl-3,4-dimethyl-4-(3-(1-hydroxy-2-methylamino)ethylphenyl)piperidine

[0107] To a solution of 1-hexanoyl-3,4-dimethyl-4-(3-[4-(4-methylphenyl)sulfonyl-4,5-dihydro-1,3-oxazol-5-yl]phenyl)piperidine (Preparation 10, 345 mg, 0.68 mmol) in anhydrous tetrahydrofuran (5 mL) at room temperature was added lithium aluminium hydride (1.0 M solution in tetrahydrofuran, 0.74 mL, 0.74 mmol) dropwise over five minutes. The solution was stirred at room temperature under a nitrogen atmosphere for 2 hours and then cooled to 0°C. The reaction was quenched cautiously by the addition of aqueous sodium hydroxide solution (1.0 mL, 1.0 M) and then ethyl acetate (20 mL) and solid sodium hydrogencarbonate (excess) were added. The mixture was stirred vigorously for 30 minutes and then filtered through Celite®, washing with ethyl acetate. The filtrate was concentrated *in vacuo* and the residue was purified by column chromatography on silica gel eluted with neat ethyl acetate and then ethyl acetate.

methanol. 0.880 ammonia (70:30 1) to give the title compound as a clear gum (120 mg)
 NMR (CDCl₃, selected data) 0.75 (d, 3H), 0.9 (t, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H),
 2.2-2.6 (m, 9H), 2.65-2.85 (m, 3H), 4.75 (m, 1H), 7.1-7.35 (m, 4H)
 MS (thermospray) M/Z (MH⁺) 347.3 C₂₂H₃₈N₂O + H requires 347.3

Preparation of Starting Materials

Preparation 1: 1-Hexyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxy phenyl)piperidine

[0108] To a solution of 1-hexyl-3,4-dimethyl-4-(3-hydroxyphenyl)piperidine (3.5 g, 12 mmol, *J. Med. Chem.*, 1993, 36, 2833) in dichloromethane (15 mL) was added triethylamine (3 mL) followed by *N*-phenyltrifluoromethanesulfonimide (6.1 g, 18 mmol) portionwise. The reaction mixture was stirred under nitrogen at room temperature for 18 hours then it was washed with aqueous sodium hydroxide solution (60 mL of 2 M). The separated aqueous layer was back-washed with dichloromethane (2 x 30 mL), after which the combined organics were dried (Na₂SO₄) and the solvent removed *in vacuo* to give a yellow oil. This was purified by column chromatography on silica gel (150 g) eluted with hexane ethyl acetate 0.880 ammonia (66:33 1) to give the title compound as a yellow oil (4.22 g).
 NMR (CDCl₃) 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.7 (m, 3H), 2.0 (m, 1H), 2.2-2.45 (m, 4H), 2.45-2.65 (m, 2H), 2.8 (m, 1H), 7.1 (d, 1H), 7.15 (s, 1H), 7.25-7.45 (m, 2H)
 MS (thermospray) M/Z (MH⁺) 422.3, C₂₀H₃₀F₃NO₃S + H requires 422.2

Preparation 2: 1-Benzyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine

[0109] To a stirred solution of 1-benzyl-3,4-dimethyl-4-(3-hydroxyphenyl)piperidine (10.16 g, 34.4 mmol, *J. Med. Chem.*, 1993, 36, 2833) in anhydrous dichloromethane (100 mL) was added triethylamine (8 mL) and the resulting solution was de-oxygenated by evacuating and flushing with nitrogen three times. *N*-Phenyltrifluoromethanesulfonimide (18.43 g, 51.6 mmol) was added and the mixture was de-oxygenated again, using the same procedure as before, and stirred overnight at room temperature under nitrogen. The reaction mixture was then diluted with dichloromethane (200 mL) and washed with aqueous sodium hydroxide solution (200 mL of 1 M). The aqueous phase was back-washed with dichloromethane (2 x 100 mL). The combined organics were dried (Na₂SO₄) and concentrated *in vacuo* to give an orange oil (ca 20 g) which was purified by column chromatography on silica gel (700 g) eluted with a gradient of ethyl acetate-hexane 0.880 ammonia (10:90 1 to 10:90 1). This gave the title compound as a colourless oil (13.98 g).
 NMR (CDCl₃) 0.75 (d, 3H), 1.35 (s, 3H), 1.55 (m, 1H), 1.95 (m, 1H), 2.25-2.5 (m, 2H), 2.5-2.65 (m, 2H), 2.85 (m, 1H), 3.45 (d, 1H), 3.6 (d, 1H), 7.1 (d, 1H), 7.15 (s, 1H), 7.2-7.45 (m, 7H)
 MS (thermospray) M/Z (MH⁺) 428.0, C₂₁H₂₄F₃NO₃S + H requires 428.2

Preparation 3: 3,4-Dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine

[0110] To a solution of 1-benzyl-3,4-dimethyl-4-(3-(*N*-ethyl)amidophenyl)piperidine (Example 16, 800 mg, 2.3 mmol) in methanol (40 mL) was added palladium on activated carbon (150 mg, Degussa type E101 NEW, Pd 10% dry weight, ca 50% water). The resulting suspension was stirred at room temperature under an atmosphere of hydrogen at 415 kPa for 1½ days. It was then filtered through Celite® to remove the catalyst residues and concentrated *in vacuo* to give a foam (610 mg). Purification by column chromatography on silica gel (30 g) eluted with CH₂Cl₂-EtOH 0.880 ammonia (50:8 1) gave the title compound as a thick gum (557 mg).
 NMR (CDCl₃, selected data) 0.7 (d, 3H), 1.25 (t, 3H), 1.4 (s, 3H), 1.95 (m, 1H), 2.15 (m, 1H), 2.75 (m, 1H), 2.95-3.15 (m, 2H), 3.25 (m, 1H), 3.5 (m, 2H), 7.3-7.45 (m, 2H), 7.5 (d, 1H), 7.7 (s, 1H)
 MS (APCI) M/Z (MH⁺) 261.5, C₁₆H₂₄N₂O + H requires 261.2

Preparation 4: 1-Hexyl-3,4-dimethyl-4-{3-[2-(trimethylsilyl)ethynyl]phenyl}piperidine

[0111] To a solution of 1-hexyl-3,4-dimethyl-4-(3-trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 1, 350 mg, 0.83 mmol) in tetrahydrofuran (12 mL) was added diisopropylamine (4 mL) and trimethylsilyl ethyne (4.5 g, 46 mmol) and the mixture was de-oxygenated by evacuating and flushing with nitrogen five times. Copper(I) iodide (6.2 mg, 0.033 mmol), and then catalytic quantities of palladium(II) acetate and 1,1'-bis(diphenylphosphino)ferrocene were added. The reaction mixture was heated to reflux under nitrogen for 8 hours, before being allowed to cool to room temperature. Water (10 mL) and dichloromethane (10 mL) were added, the phases separated and the aqueous layer further extracted with dichloromethane (2 x 10 mL). The combined organics were then dried (Na₂SO₄) and the solvent removed *in vacuo*. The residual brown oil was purified by column chromatography on silica gel (25 g) eluted with a gradient of ethyl acetate-hexane 0.880 ammonia (20:79 1 to 50:49 1) to give the title compound as an oil (150 mg).

NMR (CDCl₃) : 0.25 (s, 9H), 0.75 (d, 3H), 0.9 (m, 3H), 1.2-1.4 (m, 9H), 1.4-1.55 (m, 2H), 1.6 (m, 1H), 2.0 (m, 1H), 2.2-2.4 (m, 4H), 2.4-2.6 (m, 2H), 2.8 (m, 1H), 7.2-7.35 (m, 3H), 7.4 (s, 1H)
 MS (thermospray) : M/Z (MH⁺) 370.4; C₂₄H₃₉NSi + H requires 370.3

5 **Preparation 5: 1-Hexanoyl-3,4-dimethyl-4-(3-hydroxyphenyl)piperidine**

[0112] To a stirred solution of 3,4-dimethyl-4-(3-hydroxyphenyl)piperidine (3.8 g, 18.6 mmol, *J. Org. Chem.*, 1991, 56, 1660) in dichloromethane (30 mL) at 0°C was added triethylamine (3.9 mL, 27.8 mmol) followed by the dropwise addition of hexanoic anhydride (4.7 mL, 20.4 mmol) over 5 minutes. The reaction was stirred under a nitrogen atmosphere for 3 hours at room temperature and then quenched by the addition of saturated aqueous sodium hydrogencarbonate (50 mL). The two layers were separated and the aqueous layer was extracted with dichloromethane (3 x 50 mL). The combined organic layers were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with ethyl acetate : hexane (1 : 1). The title compound was obtained as a clear oil (4.5 g).

15 NMR (CDCl₃, selected data from a 13:9 mixture of rotamers) : 0.65 (d, 3H), 0.9 (m, 3H), 1.25-1.45 (m, 7H), 1.55-1.75 (m, 3H), 2.05 (m, 1H), 2.15 (m, 1H), 2.25-2.55 (m, 2H), 2.95 (m, 0.59H), 3.15 (m, 0.41H), 3.35 (m, 0.41H), 3.5-3.7 (m, 1.18H), 3.85 (m, 0.41H), 4.4 (m, 0.41H), 4.75 (m, 0.59H), 6.7 (d, 1H), 6.75-6.85 (m, 2H), 7.15 (t, 1H).
 MS (thermospray) : M/Z (MH⁺) 304.1; C₁₉H₂₉NO₂ + H requires 304.2

20 **Preparation 6: 1-Hexanoyl-3,4-dimethyl-4-(trifluoromethanesulfonyloxyphenyl)piperidine**

[0113] To a stirred solution of 1-hexanoyl-3,4-dimethyl-4-(3-hydroxyphenyl)piperidine (Preparation 5, 3.1 g, 10.1 mmol) in dichloromethane (30 mL) at room temperature was added triethylamine (2.82 mL, 20.2 mmol) followed by *N*-phenyltrifluoromethanesulfonimide (3.6 g, 15.1 mmol) portionwise. The reaction was stirred under a nitrogen atmosphere at room temperature for 16 hours and then aqueous sodium hydroxide (30 mL of 2 M) was added. The bi-phasic mixture was stirred vigorously for 2 hours before the two layers were separated and the aqueous layer extracted with dichloromethane (3 x 20 mL). The combined organic layers were dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with a gradient of ethyl acetate : hexane (1:2 and then 2:1). The title compound was obtained as a clear oil (3.6 g).

30 NMR (CDCl₃, selected data from a 7:5 mixture of rotamers) : 0.55-0.65 (m, 3H), 0.85-0.95 (m, 3H), 1.25-1.4 (m, 4H), 1.45 (s, 3H), 1.55-1.75 (m, 3H), 2.0-2.5 (m, 4H), 2.9 (m, 0.58H), 3.15 (m, 0.42H), 3.35 (m, 0.42H), 3.6 (m, 1.16H), 3.9 (m, 0.42H), 4.4 (m, 0.42H), 4.75 (m, 0.58H), 7.05-7.15 (m, 2H), 7.3 (m, 1H), 7.4 (m, 1H).
 MS (thermospray) : M/Z (MH⁺) 436.4; C₂₀H₂₈F₃NO₄S + H requires 436.2

35 **Preparation 7: 1-Hexanoyl-3,4-dimethyl-4-(3-methoxycarbonylphenyl)piperidine**

[0114] To a solution of 1-hexanoyl-3,4-dimethyl-4-(trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 6, 267 mg, 0.48 mmol) in anhydrous *N,N*-dimethylformamide (2 mL) was added triethylamine (0.18 mL) and methanol (0.4 mL). The mixture was de-oxygenated by evacuating and flushing with nitrogen five times. Palladium(II) acetate (4.4 mg) and 1,1'-bis(diphenylphosphino)ferrocene (8 mg) were added and the solution was purged with carbon monoxide. The reaction mixture was heated to 60°C under an atmosphere of carbon monoxide for 7 hours then it was cooled to room temperature and diluted with saturated aqueous sodium chloride solution (10 mL). The phases were separated and the aqueous layer was extracted with dichloromethane (4 x 15 mL). The combined organics were dried (Na₂SO₄) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (50 g) eluted with hexane : ethyl acetate : 0.880 ammonia (66:33:1). The title compound was obtained as a pale yellow oil (110 mg).

45 NMR (CDCl₃, selected data from a 9:7 mixture of rotamers) : 0.55-0.7 (m, 3H), 0.85-0.95 (m, 3H), 1.25-1.4 (m, 4H), 1.45 (s, 3H), 1.6-1.8 (m, 3H), 2.05-2.45 (m, 4H), 2.9 (m, 0.56H), 3.15 (m, 0.44H), 3.4 (m, 0.44H), 3.6 (m, 1.12H), 3.9 (m, 0.44H), 3.95 (s, 3H), 4.4 (0.44H), 4.7 (m, 0.56H), 7.35-7.5 (m, 2H), 7.9 (m, 1H), 7.95 (m, 1H).
 MS (thermospray) : M/Z (MH⁺) 346.3; C₂₁H₃₁NO₃ + H requires 346.2

50 **Preparation 8: 1-Hexanoyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine**

[0115] To a stirred solution of 1-hexanoyl-3,4-dimethyl-4-(trifluoromethanesulfonyloxyphenyl)piperidine (Preparation 6, 3.0 g, 6.90 mmol) in tetrahydrofuran (30 mL) at room temperature were added sequentially vinyltributyltin (2.12 mL, 7.24 mmol), lithium chloride (585 mg, 13.8 mmol), and tetrakis(triphenylphosphine)palladium(0) (80 mg, 0.69 mmol). The mixture was heated to reflux under a nitrogen atmosphere for 1½ hours at which time a few crystals of 4-*tert*-butylcatechol were added. Heating at reflux was then continued for a further 16 hours. The mixture was cooled and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with a gradient of ethyl

acetate hexane (1:10 to 1:3). The title compound was obtained as a clear oil (2.1 g).
 NMR (CDCl₃, selected data from a 5:4 mixture of rotamers): 0.55-0.7 (m, 3H), 0.85-1.0 (m, 3H), 1.25-1.4 (m, 4H), 1.4 (s, 3H), 1.6-1.75 (m, 3H), 2.05-2.45 (m, 4H), 2.9 (m, 0.56H), 3.15 (m, 0.44H), 3.35 (m, 0.44H), 3.6 (m, 1.12H), 3.9 (m, 0.44H), 4.4 (m, 0.44H), 4.7 (m, 0.56H), 5.25 (d, 1H), 5.75 (d, 1H), 6.7 (dd, 1H), 7.15 (m, 1H), 7.2-7.35 (m, 3H).
 MS (APCI) M/Z (MH⁺) 314.5. C₂₁H₃₁NO + H requires 314.2.

Preparation 9: 1-Hexanoyl-3,4-dimethyl-4-(3-formylphenyl)piperidine

[0116] To a solution of 1-hexanoyl-3,4-dimethyl-4-(3-vinylphenyl)piperidine (Preparation 8, 2.4 g, 7.67 mmol) in acetone (20 mL) at room temperature was added water (5 mL), 4-methylmorpholine N-oxide (1.1 g, 9.20 mmol) and finally osmium tetroxide (3.83 mL, 2.5 wt % solution in *tert*-butanol). The yellow solution was stirred at room temperature for 1 hour and then sodium periodate (4.92 g, 23.0 mmol) was added in one portion. After stirring the reaction for 3 hours a heavy precipitate had developed and the reaction mixture was filtered through Celite® washing with acetone. The filtrate was concentrated *in vacuo*, the crude oil was dissolved in dichloromethane, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with ethyl acetate hexane (1:1). The title compound was isolated as clear oil (2.0 g).
 NMR (CDCl₃, selected data from a 1:1 mixture of rotamers): 0.55-0.7 (m, 3H), 0.85-0.95 (m, 3H), 1.25-1.4 (m, 4H), 1.45 (s, 3H), 1.55-1.8 (m, 3H), 2.1-2.5 (m, 4H), 2.95 (m, 0.5H), 3.15 (m, 0.5H), 3.4 (m, 0.5H), 3.6 (m, 1H), 3.9 (m, 0.5H), 4.4 (m, 0.5H), 4.75 (m, 0.5H), 7.45-7.6 (m, 2H), 7.7 (m, 1H), 7.75 (m, 1H), 10.0 (s, 1H).
 MS (thermospray) M/Z (MH⁺) 316.3. C₂₀H₂₉NO₂ + H requires 316.2.

Preparation 10: 1-Hexanoyl-3,4-dimethyl-4-{3-[4-(4-methylphenyl)sulfonyl-4,5-dihydro-1,3-oxazol-5-yl]phenyl}piperidine


[0117] To a solution of 1-hexanoyl-3,4-dimethyl-4-(3-formylphenyl)piperidine (Preparation 9, 758 mg, 2.40 mmol) in ethanol (20 mL) was added [(4-methylphenyl)sulfonyl]methyl isocyanide (460 mg, 2.34 mmol) followed by sodium cyanide (12 mg, 0.24 mmol). The mixture was stirred at room temperature under a nitrogen atmosphere for five hours and then concentrated *in vacuo*. The residue was purified by column chromatography on silica gel using a gradient elution of hexane ethyl acetate (67:33 to 0:100). The title compound was isolated as a clear oil (909 mg).
 NMR (CDCl₃, selected data from a 1:1 mixture of rotamers): 0.55-0.65 (m, 3H), 0.85-0.95 (m, 3H), 1.25-1.45 (m, 7H), 1.55-1.75 (m, 3H), 2.05-2.45 (m, 4H), 2.45 (s, 3H), 2.9 (m, 0.5H), 3.15 (m, 0.5H), 3.35 (m, 0.5H), 3.6 (m, 1H), 3.9 (m, 0.5H), 4.4 (m, 0.5H), 4.7 (m, 0.5H), 5.0 (d, 1H), 6.05 (d, 1H), 7.1-7.3 (m, 4H), 7.3-7.45 (m, 3H), 7.85 (d, 2H).
 MS (thermospray) M/Z (MH⁺) 511.1. C₂₉H₃₈N₂O₄S + H requires 511.3.

Biological Activity

[0118] The K_i values of certain compounds of the present invention in the opioid receptor binding assays were determined, and the compounds of Examples 4, 8, 18 and 20 were all found to have K_i values of 4000 nM or less for the μ receptor. The compounds of the invention also possess affinity at the δ and κ opioid receptors.

Claims

1. A compound of formula I



wherein

A represents a single bond, C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH;

D represents H, OH, CN, $N(R^4)(R^5)$, $N(H)R^6$, $C(O)N(R^4)(R^5)$, $C(O)OR^7$, $C(O)R^8$, $C(=NR^{9a})R^8$, or $C(=NOR^{9b})R^8$;

provided that when A represents C_{2-4} alkenylene or C_{2-4} alkynylene, and D represents OH, $N(R^4)(R^5)$ or $N(H)R^6$, then D is not directly attached to an unsaturated carbon atom;

and provided that when A represents a single bond, then D does not represent H, OH, $N(R^4)(R^5)$ or $N(H)R^6$; R^4 and R^5 independently represent H, C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), or R^4 and R^5 , together with the N-atom to which they are attached, form a 4- to 7-membered heterocyclic ring, which ring optionally contains one or more additional heteroatoms selected from oxygen, nitrogen and sulfur and which ring is optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, OH, =O, nitro, amino or halo, R^6 represents $C(O)R^{10a}$, $C(O)OR^{10b}$ or $S(O)_2R^{10c}$;

R^{10a} to R^{10c} independently represent C_{1-4} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl (which four groups are all optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms)), or R^{10a} represents H;

R^7 and R^8 independently represent H, C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl or C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms);

R^{9a} and R^{9b} independently represent C_{1-6} alkyl, C_{3-8} cycloalkyl, aryl, C_{1-4} alkylphenyl, which latter four groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), or R^{9b} represents H;

R^1 and R^2 are each independently H or C_{1-4} alkyl;

R^3 represents aryl (optionally substituted by one or more substituents selected from OH, nitro, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms) and $-N(R^{11a})(R^{11b})$), C_{1-10} alkyl, C_{3-10} alkenyl or C_{3-10} alkynyl wherein said alkyl, alkenyl or alkynyl groups are optionally substituted and/or terminated by one or more substituents selected from OR^{11c} , $S(O)_pR^{11d}$, CN, halo, C_{1-6} alkoxy carbonyl, C_{2-6} alkanoyl, C_{2-6} alkanoyloxy, C_{3-8} cycloalkyl, C_{4-9} cycloalkanoyl, $N(R^{12a})S(O)_2R^{13}$, Hal^1 , aryl, adamantyl (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , $CONH_2$, C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), or $-W-A^1-N(R^{12b})(R^{12c})$;

p is 0, 1 or 2;

W represents a single bond, $C(O)$ or $S(O)_q$;

A^1 represents a single bond or C_{1-10} alkylene;

provided that when both W and A^1 represent single bonds, then the group $-N(R^{12b})(R^{12c})$ is not directly attached to an unsaturated carbon atom; q is 0, 1 or 2;

R^{11a} to R^{11d} each independently represent H, C_{1-10} alkyl, C_{3-10} alkenyl, C_{3-10} alkynyl, C_{3-8} cycloalkyl, C_{1-4} alkylphenyl, aryl (which latter six groups are optionally substituted by one or more substituents selected

from OH, nitro, amino, halo, CN, CH₂CN, CONH₂, C₁₋₄ alkyl, C₁₋₄ alkoxy and C₁₋₅ alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)) or Het² provided that R^{11d} does not represent H when p represents 1 or 2.

R^{12a} to R^{12c} each independently represent H, C₁₋₁₀ alkyl, C₃₋₁₀ alkenyl, C₃₋₁₀ alkynyl, C₃₋₈ cycloalkyl, C₁₋₄ alkylphenyl, aryl (which latter six groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH₂CN, CONH₂, C₁₋₄ alkyl, C₁₋₄ alkoxy and C₁₋₅ alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), Het³, or R^{12b} and R^{12c} together represent unbranched C₂₋₆ alkylene which alkylene group is optionally interrupted by O, S and/or an N(R¹⁴) group and is optionally substituted by one or more C₁₋₄ alkyl groups;

R¹³ represents C₁₋₆ alkyl, C₃₋₈ cycloalkyl, C₁₋₄ alkylphenyl or aryl, which four groups are optionally substituted by one or more substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, OH, nitro, amino or halo.

R¹⁴ represents H, C₁₋₆ alkyl, C₃₋₈ cycloalkyl, A²-(C₃₋₈ cycloalkyl) or A²-aryl,

A² represents C₁₋₆ alkylene.

Het¹, Het² and Het³ independently represent 3- to 8-membered heterocyclic groups, which groups contain at least one heteroatom selected from oxygen, sulfur and/or nitrogen, which groups are optionally fused to a benzene ring, and which groups are optionally substituted in the heterocyclic and/or fused benzene ring part by one or more substituents selected from OH, =O, nitro, amino, halo, CN, aryl, C₁₋₄ alkyl, C₁₋₄ alkoxy and C₁₋₅ alkanoyl (which latter three groups are optionally substituted by one or more halo atoms),

X is H, halo, C₁₋₄ alkyl or C₁₋₄ alkoxy (which latter two groups are optionally substituted by one or more halo atoms)


n is 0, 1 or 2

or pharmaceutically or veterinarily acceptable derivatives thereof

2. A compound as claimed in Claim 1 wherein the group A-D is attached in the *meta*-position relative to the piperidine ring
3. A compound as claimed in Claim 1 or Claim 2 wherein R¹ represents C₁₋₂ alkyl
4. A compound as claimed in any one of Claims 1 to 3 wherein R² represents H or C₁₋₂ alkyl
5. A compound as claimed in any one of Claims 1 to 4 wherein R³ represents saturated C₁₋₁₀ alkyl optionally substituted by one or more substituents selected from OR^{11c}, CN, halo, C₂₋₄ alkanoyl, C₁₋₄ alkoxy carbonyl, N(R^{12a})SO₂R¹³, Het¹, aryl (which latter group is optionally substituted by one or more substituents selected from OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₅ alkanoyl, halo, nitro, amino, CN and CONH₂), or -W-A¹-N(R^{12b})(R^{12c})
6. A compound as claimed in any one of Claims 1 to 5 wherein R^{11c} represents H, C₁₋₆ alkyl or aryl (which latter groups is optionally substituted by one or more substituents selected from OH, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₂₋₅ alkanoyl, halo, nitro, amino, CN and CONH₂), R^{12a} to R^{12c} independently represent H, C₁₋₄ alkyl, C₁₋₂ alkylphenyl or aryl (which latter three groups are optionally substituted by one or more substituents selected from halo, C₁₋₄ alkyl or C₁₋₄ alkoxy), W represents C(O), and/or A¹ represents a single bond
7. A compound as claimed in any one of Claims 1 to 6 wherein R¹³ represents C₁₋₄ alkyl, C₁₋₂ alkylphenyl or aryl (which three groups are all optionally substituted by one or more substituents selected from halo, C₁₋₄ alkyl or C₁₋₄ alkoxy)
8. A compound as claimed in any one of Claims 1 to 7 wherein A represents a single bond, C₁₋₄ alkylene, C₂₋₄ alkenylene or C₂₋₄ alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more OH and/or methyl groups
9. A compound as claimed in any one of Claims 1 to 8 wherein D represents H, OH, CN, N(H)R⁴, N(H)C(O)R^{10a}, N(H)C(O)OR^{10b}, N(H)S(O)₂R^{10c}, C(O)N(R⁴)(R⁵), C(O)OR⁷, C(O)R⁸ or C(=NOH)R⁸, R⁴ and R⁵ independently represent H, C₁₋₄ alkyl or C₁₋₃ alkylphenyl, which latter two groups are optionally substituted by C₁₋₄ alkoxy, R⁷ and R⁸ independently represent H or C₁₋₄ alkyl, and/or R^{10a} to R^{10c} independently represent C₁₋₄ alkyl, which group is optionally substituted by one or more halo atoms
10. A compound as claimed in any one of Claims 1 to 9 wherein R³ represents saturated C₁₋₇ alkyl, optionally substituted by one or more substituents selected from CN, O-(C₁₋₆ alkyl), phenyl, or O-(phenyl)

11. A compound as claimed in any one of Claims 1 to 10 wherein X represents halo
12. A compound as claimed in any one of Claims 1 to 11 wherein n represents 0 or 1.
13. A compound as defined in any one of Claims 1 to 12, for use as a medicament.
14. A compound as defined in any one of Claims 1 to 12, for use as an animal medicament.
15. A formulation comprising a compound as defined in any one of Claims 1 to 12, in admixture with a pharmaceutically, or a veterinary, acceptable adjuvant, diluent or carrier
16. A formulation as claimed in Claim 15, which is a veterinary formulation
17. The use of a compound as defined in any one of Claims 1 to 12, in the manufacture of a medicament for the curative or prophylactic treatment of a disease mediated via an opiate receptor.
18. The use as claimed in Claim 17, wherein the disease is pruritus.
19. A method of treating or preventing a disease mediated by an opiate receptor, which comprises administering a therapeutically effective amount of a compound as defined in any one of Claims 1 to 12, to a patient in need of such treatment
20. A process for the preparation of a compound as defined in Claim 1, which comprises:

a) for compounds of formula I in which A represents C_{2-4} alkynylene (in which group the carbon-carbon triple bond is α, β to the benzene ring), which alkynylene group is optionally substituted at the 3- and/or the 4-C (relative to the benzene ring) by one or more substituents defined in Claim 1 in respect of A, and/or one of the groups defined in Claim 1 in respect of D, or (when D is not attached at the 3- or 4-C) which alkynylene group is substituted at the 2-C (relative to the benzene ring) by CN, $C(O)N(R^4)(R^5)$, $C(O)OR^7$, $C(O)R^8$, $C(=NR^{9a})R^8$, or $C(=NOR^{9b})R^8$, reaction of a corresponding compound of formula II,




wherein L^1 is a leaving group, and R^1 , R^2 , R^3 , X and n are as defined in Claim 1, with a compound of formula III,



wherein M represents (as appropriate) H, a tin-containing moiety, a boron derivative, a zinc halide, a magnesium halide or an alkali metal, A^3 represents a single bond or C_{1-2} alkylene (optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, OH or halo), and D is as defined in Claim 1, provided that when A^3 represents a single bond, then D does not represent H, OH, $N(R^4)(R^5)$ or $N(H)R^6$, wherein R^4 , R^5 and R^6 are as defined in Claim 1.

b) for compounds of formula I in which A represents C_{2-4} alkenylene (in which group the carbon-carbon double bond is α, β to the benzene ring), which alkenylene group is optionally substituted at the 2-C (relative to the


benzene ring) by C₁₋₄ alkyl and also optionally substituted at the 3- and/or 4-C (relative to the benzene ring) by one or more of the substituents defined in Claim 1 in respect of A and/or one of the groups defined in Claim 1 in respect of D, or which alkenylene group is substituted at the 2-C (relative to the benzene ring) by CN, C(O)N(R⁴)(R⁵), C(O)OR⁷, C(O)R⁸, C(=NR^{9a})R⁸, or C(=NOR^{9b})R⁸. reaction of a corresponding compound of formula II, as defined above, with a compound of formula IV,



IV

wherein the dashed bond represent optional *cis*- or *trans*- geometry R¹⁵ represents H or C₁₋₄ alkyl, A³ and M are as defined above, and D is as defined in Claim 1,

c) for compounds of formula I in which A represents a single bond and D represents CN, reaction of a compound of formula V




V

wherein R¹ R² R³ X and n are as defined in Claim 1, with an alkali metal cyanide.

d) for compounds of formula I in which A represents C₁₋₄ alkylene, C₂₋₄ alkenylene or C₂₋₄ alkynylene, which alkylene alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C₁₋₄ alkyl C₁₋₄ alkoxy halo or OH, and D represents NH₂ (which is attached to a CH₂ group), reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C₁₋₃ alkylene, C₂₋₃ alkenylene or C₂₋₃ alkynylene, which alkylene alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C₁₋₄ alkyl, C₁₋₄ alkoxy, halo or OH, and D represents CN

e) for compounds of formula I in which D represents C(O)NH₂, controlled hydrolysis of a corresponding compound of formula I in which D represents CN,

f) for compounds of formula I in which A represents a single bond and D represents C(O)-(C₁₋₆ alkyl) or C(O)-(C₁₋₄ alkylphenyl) which alkyl and alkylphenyl groups are both optionally substituted by one or more of the substituents defined in Claim 1 in respect of R⁸, hydrolysis of a corresponding compound of formula IX,



IX

wherein R^{15} represents C_{1-6} alkyl, R^{16} represents H, C_{1-5} alkyl, phenyl or C_{1-3} alkylphenyl which latter three groups are all optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), the dashed bond indicates optional *cis*- or *trans*- geometry, and R^1 , R^2 , R^3 , X and n are as defined in Claim 1,

g) for compounds of formula I in which D represents $C(O)R^8$, wherein R^8 is as defined in Claim 1 provided that it does not represent H, reaction of a corresponding compound of formula I in which D represents CN with an organometallic compound capable of delivering an R^{8a} -containing anion, wherein R^{8a} is defined as for R^8 in Claim 1 provided that it does not represent H,


h) for compounds of formula I in which A represents a single bond and D represents $C(O)OR^7$, wherein R^7 is as defined in Claim 1 provided that it does not represent H, reaction of a corresponding compound of formula V as defined above, with carbon monoxide and an alcohol of formula $R^{7a}OH$, wherein R^{7a} is defined as for R^7 in Claim 1 provided that it does not represent H;

i) for compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents OH (which is attached to a CH_2 group), reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $C(O)OR^{7a}$, wherein R^{7a} is as defined above,

j) for compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are *gem*-disubstituted with two C_{1-4} alkyl groups (a to D) and are optionally substituted by one or more further substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents OH, reaction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $C(O)OR^{7a}$, wherein R^{7a} is as defined above, with a C_{1-4} alkyl-delivering organometallic compound,

k) for compounds of formula I in which D represents $C(O)N(R^4)(R^5)$, wherein R^4 and R^5 are as defined in Claim 1.

A) reaction of a corresponding compound of formula I in which D represents $C(O)OR^{7a}$, wherein R^{7a} is as defined above, with a compound of formula XI,



XI

or an acid addition salt thereof, wherein R^4 and R^5 are as defined in Claim 1,

B) reaction of a corresponding compound of formula I in which D represents $C(O)OH$ with a compound of formula XI, as defined above:

l) for compounds of formula I in which D represents $C(O)OH$, hydrolysis of a corresponding compound of

formula I in which D represents $C(O)OR^{7a}$ wherein R^{7a} is as defined above;

m) for compounds of formula I in which D represents $N(H)R^6$, wherein R^6 is as defined in Claim 1, reaction of a corresponding compound of formula I in which D represents NH_2 with a compound of formula XII,



XII

wherein R^6 is as defined in Claim 1 and L^1 is as defined above.


n) for compounds of formula I in which A represents C_{1-4} alkyl and D represents $N(R^4)(R^5)$ or $N(H)C(O)R^{10a}$ attached at the 1-, 2- or 3-C (relative to the benzene ring), wherein R^4 , R^5 and R^{10a} are as defined in Claim 1, reaction of a corresponding compound of formula I in which A represents C_{1-4} alkenylene unsaturated α , β -, β,γ - or γ,δ - (respectively) relative to the benzene ring and D represents H with a compound of formula XII as defined above or a compound of formula XIII,



XIII

wherein R^{10a} is as defined in Claim 1,

o) for compounds of formula I in which A represents C_{2-4} alkylene optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents OH, oxidation of a corresponding borane adduct of formula XIV



XIV

wherein x is 1, 2 or 3, y is (as appropriate) (3-x) or 1, R^{17} is (as appropriate) H, halo, an alkyl or a cycloalkyl group providing one or two bonds to boron, A represents (as appropriate) C_{2-4} alkylene optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and R^1 , R^2 , R^3 , X and n are as defined in Claim 1.

p) for compounds of formula I in which A represents a C_{2-4} alkylene group substituted (α to D) with an OH group and D represents OH, reaction of a corresponding compound of formula I in which A represents a C_{2-4} alkenylene group and D represents H with a dihydroxylating reagent,


q) for compounds of formula I in which A represents a single bond or a C_{1-2} alkylene group (as appropriate) and D represents $C(O)H$, reaction of a corresponding compound of formula I in which A represents a C_{2-4} alkylene group substituted (α to D) with an OH group and D represents OH with a reagent that effects 1,2-diol oxidative cleavage,

r) for compounds of formula I in which D represents $C(=NR^{9a})R^8$ or $C(=NOR^{9b})R^8$, wherein R^8 , R^{9a} and R^{9b} are as defined in Claim 1, reaction of a corresponding compound of formula I in which D represents $C(O)R^8$ with a compound of formula XV,



XV


or a compound of formula XVI,



XVI

wherein R^{9a} and R^{9b} are as defined in Claim 1;


s) for compounds of formula I in which A represents C_{1-4} alkylene substituted (α to D) with an OH group and D represents N(H)CH_3 (at the alkylene chain terminus), reduction of a corresponding compound of formula XVII,



XVII

wherein r is 0, 1 or 2, L^2 represents H or a group capable, when attached to a C_2 alkylene unit, of undergoing 1,2-elimination (relative to L^2), and R^1 , R^2 , R^3 , X and n are as defined in Claim 1;


t) for compounds of formula I wherein R^3 represents C_1 alkyl optionally substituted by C_{3-8} cycloalkyl, Het^1 , aryl, adamantyl, (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , CONH_2 , C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), or R^3 represents C_{2-10} alkyl, C_{3-10} alkenyl or C_{3-10} alkynyl (which three groups are all optionally substituted by one or more of the relevant substituents identified in Claim 1 in respect to R^3), which alkyl, alkenyl or alkynyl groups are attached to the piperidine nitrogen atom via a CH_2 group, wherein Het^1 is as defined in Claim 1, reduction of a corresponding compound of formula XIX,



XIX

wherein R^{31} represents H, C_{3-8} cycloalkyl, Het^1 , aryl, adamantyl, (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , CONH_2 , C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), C_{1-9} alkyl, C_{2-9} alkenyl or C_{2-9} alkynyl, which alkyl, alkenyl or alkynyl groups are optionally substituted and/or terminated by one or more substituents selected from OR^{11c} , $\text{S(O)}_p\text{R}^{11d}$, CN, halo, C_{1-6} alkoxy carbonyl, C_{2-6} alkanoyl, C_{2-6} alkanoyloxy, C_{3-8} cycloalkyl, C_{4-9} cycloalkanoyl, $\text{N(R}^{12a})\text{S(O)}_2\text{R}^{13}$, Het^1 , aryl, adamantyl (which latter two groups are optionally substituted by one or more substituents selected from OH, nitro, amino, halo, CN, CH_2CN , CONH_2 , C_{1-4} alkyl, C_{1-4} alkoxy and C_{1-5} alkanoyl (which latter three groups are optionally substituted by one or more halo atoms)), or $-\text{W-A}^1-\text{N(R}^{12b})\text{(R}^{12c})$, and R^1 , R^2 , R^{11c} , R^{11d} , R^{12a} to R^{12c} , R^{13} , Het^1 ,

n, p, W, X, A¹, A and D are as defined in Claim 1.
 u) reaction of a corresponding compound of formula XX,




wherein R¹, R², A, D, X and n are as defined in Claim 1, with a compound of formula VIII,



wherein R³ is as defined in Claim 1 and L¹ is as defined above

v) for compounds of formula I wherein R³ represents C₁ alkyl, which, in place of being optionally substituted by the substituents as defined in Claim 1, is instead optionally substituted by R³¹, wherein R³¹ is as defined above, reaction of a corresponding compound of formula XX, as defined above, with a compound of formula XXII

25



wherein R³¹ is as defined above, in the presence of a reducing agent,

w) for compounds of formula I wherein R³ is a C₁₋₁₀ alkyl, C₄₋₁₀ alkenyl or C₄₋₁₀ alkynyl group that is fully saturated from 1- to 3-C (relative to the piperidine N-atom), and which R³ group is substituted at 2-C (relative to the piperidine N-atom) by S(O)R^{11d}, S(O)₂R^{11d}, alkanoyl, cycloalkanoyl, alkoxy carbonyl, CN, -C(O)-A¹-N(R^{12b})(R^{12c}), -S(O)-A¹-N(R^{12b})(R^{12c}) or -S(O)₂-A¹-N(R^{12b})(R^{12c}), wherein R^{11d}, R^{12b}, R^{12c} and A¹ are as defined in Claim 1, reaction of a corresponding compound of formula XX, as defined above, with a compound of formula XXIII,

35




wherein R^{3a} represents R³ as defined in Claim 1 except that it does not represent aryl, and that the R^{3a} chain contains an additional carbon-carbon double bond α,β to the Z-substituent, and Z represents S(O)R^{11d}, S(O)₂R^{11d}, alkanoyl, cycloalkanoyl, alkoxy carbonyl, CN, -C(O)-A¹-N(R^{12b})(R^{12c}), -S(O)-A¹-N(R^{12b})(R^{12c}) or -S(O)₂-A¹-N(R^{12b})(R^{12c}), wherein R^{11d}, R^{12b}, R^{12c} and A¹ are as defined in Claim 1.

45

x) for compounds of formula I in which A represents C₂₋₄ alkylene substituted (α to D) with an OH group and D represents N(R⁴)(R⁵) (at the alkylene chain terminus), and R⁴ and R⁵ are as defined in Claim 1, reaction of a compound of formula XXIV


50

55




wherein R^1 , R^2 , R^3 , X and n are as defined in Claim 1 and r is as defined above, with a compound of formula XI, as defined above

y) for compounds of formula I in which D represents $N(H)R^4$, wherein R^4 is as defined in Claim 1 provided that it does not represent aryl, reduction of a corresponding compound of formula XXV,



wherein R^{4b} and R^{4c} , together with the carbonyl group to which they are attached, form a C_{1-6} alkanal, C_{3-6} alkanone, C_{3-8} cycloalkanone, phenyl(C_{1-4})alkanal or phenyl(C_{2-4})alkanone group, which five groups are optionally substituted by one or more substituents selected from nitro, halo, C_{1-4} alkyl or C_{1-4} alkoxy (which latter two groups are optionally substituted by one or more halo atoms), and R^1 , R^2 , R^3 , A , X and n are as defined in Claim 1 (provided that the $-N=C(R^{4b})(R^{4c})$ group is not directly attached to an unsaturated carbon atom), z) for compounds of formula I in which A represents C_{1-4} alkylene, C_{2-4} alkenylene or C_{2-4} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $N(R^4)(R^5)$ (attached to a CH_2 group) wherein R^4 and R^5 are as defined in Claim 1, reduction of a corresponding compound of formula I in which A represents (as appropriate) a single bond, C_{1-3} alkylene, C_{2-3} alkenylene or C_{2-3} alkynylene, which alkylene, alkenylene or alkynylene groups are optionally substituted by one or more substituents selected from C_{1-4} alkyl, C_{1-4} alkoxy, halo or OH, and D represents $C(O)N(R^4)(R^5)$

aa) conversion of one functional group on an alkyl, heterocyclic or aryl group in a compound of formula I to another



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 072 592 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
07.02.2001 Bulletin 2001/06

(51) Int Cl.7: **C07D 211/14, C07D 211/22,
C07D 211/32, C07D 211/34,
A61K 31/445, A61P 17/04**

(43) Date of publication A2:
31.01.2001 Bulletin 2001/05

(21) Application number: **00304219.9**

(22) Date of filing: **18.05.2000**

(84) Designated Contracting States
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **28.05.1999 GB 9912411**

(71) Applicants
• **PFIZER INC.**
New York, N.Y. 10017-5755 (US)
Designated Contracting States:
BE CH DE DK ES FI FR GR IE IT LI LU NL PT AT CY
• **Pfizer Limited**
Sandwich Kent CT13 9NJ (GB)
Designated Contracting States:
GB


(72) Inventors.
• **Armer, Richard Edward**
Newhouse, Scotland ML1 5SH (GB)
• **Gethin, David Morris**
Sandwich, Kent CT13 9NJ (GB)
• **Gibson, Stephen Paul**
Sandwich, Kent CT13 9NJ (GB)
• **Tommasini, Ivan**
Sandwich, Kent CT13 9NJ (GB)

(74) Representative: **Eddowes, Simon et al**
Urquhart-Dykes & Lord,
30 Welbeck Street
London W1G 8ER (GB)

(54) **4-arylpiperidine derivatives for the treatment of pruritus**

(57) There is provided a compound of formula I,

wherein A, D, R¹, R², R³, X and n have meanings given in the description, which are useful in the prophylaxis and in the treatment of diseases mediated by opiate receptors, such as pruritus.



EP 1 072 592 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 46 of the European Patent Convention EP 00 30 4219
shall be considered, for the purposes of subsequent
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL7)
X	WO 98 51311 A (SYNAPTIC PHARMA CORP) 19 November 1998 (1998-11-19) * example 99 *	1,2,4,5, 8,9,12	C07D211/14 C07D211/22 C07D211/32 C07D211/34
X	EP 0 494 717 A (SHELL INT RESEARCH) 15 July 1992 (1992-07-15) * examples 56-61 *	1,2,4,5, 8-10,12	A61K31/445 A61P17/04
X	US 4 292 321 A (PATTISON IAN C) 29 September 1981 (1981-09-29) * example 41 *	1,2,4,5, 8,9,12, 13	
X	EP 0 007 067 A (CIBA GEIGY AG) 23 January 1980 (1980-01-23) * example 2 *	1,2,4,5, 8-10,12, 13	
			TECHNICAL FIELDS SEARCHED (InCL7)
			C07D A61K A61P
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims</p> <p>Claims searched completely</p> <p>Claims searched incompletely:</p> <p>Claims not searched</p> <p>Reason for the limitation of the search</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		28 November 2000	De Jong, B
CATEGORY OF CITED DOCUMENTS			
<p>X particularly relevant if taken alone</p> <p>Y particularly relevant if combined with another document of the same category</p> <p>A technological background</p> <p>O non-written disclosure</p> <p>P intermediate document</p>		<p>T theory or principle underlying the invention</p> <p>E earlier patent document, but published on, or after the filing date</p> <p>D document cited in the application</p> <p>L document cited for other reasons</p> <p>... ..</p> <p>3 member of the same patent family, corresponding document</p>	

EP 00 30 4219 (1998-11-19)

European Patent
OfficeINCOMPLETE SEARCH
SHEET CApplication Number
EP 00 30 4219

Although claim 19 is directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compounds.

Claim(s) searched completely:
2,17-19

Claim(s) searched incompletely:
1,3-16,20

Reason for the limitation of the search:

Present claims 1,3-16, 20 relate to an extremely large number of possible compounds. Support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of formula I, wherein the group A-D is attached in the meta-position relative to the piperidine ring.

European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 00 30 4219

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (InCL17)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	BELLIOTTI, THOMAS R. ET AL: "Isoindolinone enantiomers having affinity for the dopamine D4 receptor" BIOORG. MED. CHEM. LETT. (1998), 8(12), 1499-1502 XP002154033 compound 10b	1,2,4,5, 8,9,12, 13	
X	WO 97 45419 A (WARNER LAMBERT CO ; WISE LAWRENCE DAVID (US); WUSTROW DAVID JUERGEN) 4 December 1997 (1997-12-04) * examples 17,36 *	1,2,4,5, 8,9,13	
X	US 5 885 994 A (KESTEN SUZANNE ROSS ET AL) 23 March 1999 (1999-03-23) * example 15 *	1,2,4,5, 8,9,13	TECHNICAL FIELDS SEARCHED (InCL17)
A	EP 0 287 339 A (LILLY CO ELI) 19 October 1988 (1988-10-19) * claims *	1,13,17, 19	
P,X	EP 0 938 898 A (PFIZER LTD ; PFIZER (US)) 1 September 1999 (1999-09-01) * claim 1 *	1-20	
P,X	WO 99 59971 A (GETHIN DAVID MORRIS ; GIBSON STEPHEN PAUL (GB); SMITH JULIAN DUNCAN) 25 November 1999 (1999-11-25) * claim 1 *	1-20	

EPO FORM 1503 03/92 (P44C18)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 4219

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-11-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9851311 A	19-11-1998	AU 7687298 A	08-12-1998
EP 0494717 A	15-07-1992	BR 9205423 A	15-03-1994
		CN 1063101 A	29-07-1992
		CS 9200044 A	15-07-1992
		WO 9212130 A	23-07-1992
		HU 65119 A	28-04-1994
		JP 6506441 T	21-07-1994
		RU 2097375 C	27-11-1997
		TR 28066 A	11-12-1995
		US 5489599 A	06-02-1996
		ZA 9200149 A	30-09-1992
US 4292321 A	29-09-1981	US 4364954 A	21-12-1982
EP 0007067 A	23-01-1980	DD 144770 A	05-11-1980
		DK 282379 A	06-01-1980
		ES 482138 A	01-04-1980
		FI 792076 A	06-01-1980
		JP 55011592 A	26-01-1980
		NO 792233 A	08-01-1980
		PL 216817 A	24-03-1980
		PT 69871 A	01-08-1979
		ZA 7903336 A	25-06-1980
WO 9745419 A	04-12-1997	AU 3072797 A	05-01-1998
		BG 102959 A	31-08-1999
		BR 9709630 A	10-08-1999
		CN 1219930 A	16-06-1999
		CZ 9803846 A	17-03-1999
		EP 0906294 A	07-04-1999
		HR 970301 A	30-04-1998
		JP 2000511534 T	05-09-2000
		NO 985531 A	27-11-1998
		PL 330195 A	26-04-1999
		SK 162798 A	11-06-1999
US 5885994 A	23-03-1999	NONE	
EP 0287339 A	19-10-1988	AT 110057 T	15-09-1994
		AU 596290 B	26-04-1990
		AU 1462488 A	20-10-1988
		CA 1321792 A	31-08-1993
		CN 88102191 A, B	02-11-1988
		DE 3851081 D	22-09-1994
		DE 3851081 T	16-02-1995

EPO FORM 1006

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 4219

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-11-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0287339 A		DK 204388 A	05-01-1989
		EG 18864 A	29-06-1995
		ES 2058265 T	01-11-1994
		HU 46892 A,B	28-12-1988
		IE 64508 B	09-08-1995
		IL 86061 A	15-07-1992
		JP 2661699 B	08-10-1997
		JP 63277661 A	15-11-1988
		KR 9615087 B	24-10-1996
		MX 11117 A	01-11-1993
		NZ 224236 A	28-08-1990
		PH 24752 A	01-10-1990
		PT 87233 A,B	01-05-1988
		SU 1598869 A	07-10-1990
		US 5422356 A	06-06-1995
		US 4891379 A	02-01-1990
		US 4992450 A	12-02-1991
		US 5064834 A	12-11-1991
		US 5319087 A	07-06-1994
		ZA 8802640 A	27-12-1989
EP 0938898 A	01-09-1999	BR 9804749 A	28-03-2000
		JP 11228411 A	24-08-1999
		US 5972962 A	26-10-1999
WO 9959971 A	25-11-1999	AU 3531299 A	06-12-1999

EPO FORM P0439

For more details about this annex, see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.